Zoological Record

Vol. 90, Sect. 5, 1953

ECHINODERMA

COMPILED BY

A. G. BRIGHTON, M.A.

LONDON
PUBLISHED BY
THE ZOOLOGICAL SOCIETY OF LONDON

PRICE TWO SHILLINGS AND SIXPENCE
June, 1955

SEPARATE SECTIONS OF THE ZOOLOGICAL RECORD

In addition to the complete bound volume, the separate Sections may be obtained singly, bound in printed paper covers.

The following are the Sections and their prices, viz .:-

							8.	d.
1.	Comprehen	sive Zo	ology		•••	 	2	6
2.	Protozoa					 	7	6
3.	Porifera					 	2	0
4.	Coelenterat	a				 	4	0
5.	Echinodern	ıa				 	2	6
6.	Vermes					 	10	0
7.	Brachiopod	a				 	3	0
8.	Bryozoa					 	2	0
9.	Mollusca		.1			 	10	0
10.	Crustacea					 	5	0
11.	Trilobita					 	-3	0
12.	Arachnida					 	7	6
*13.	Insecta					 	30	0
14.	Protochorda	ata				 	2	0
15.	Pisces					 	7	0
16A	. Amphibia					 	3	6
16в	. Reptilia					 	4	0
17.	Aves					 	7	6
18.	Mammalia					 	7	6
19.	List of New	Genera	a and S	Subgene	era	 	3	0

^{*} Obtainable only from the Commonwealth Institute of Entomology, 56, Queen's Gate, London, S.W.7.

5. ECHINODERMA

COMPILED BY

A. G. BRIGHTON, M.A.

CONTENTS

						PAG
I.	TITLES		 	 	 	1
II.	SUBJECT I	NDEX	 	 	 	12
II.	Systemati	c INDEX	 	 	 	15

I.—TITLES

1.—[Anon.] Regeneration of a starfish, Nat, Hist, notes nat, Hist, Soc. Jamaica No. 59 1953; 188 figs. 1–5.

Alfert, M. see Mazia, D.

2.—Anderson, J. M. Structure and function in the pyloric casea of *Asterias forbesi*. Biol. Bull. Woods Hole 105 1953: 47-61 pls. i-iii.

Anguéra, G. see Decourt, P.

- 3.—Bäckström, S. Studies on the animalizing action of iodosobenzoic acid in the sea urchin development. Ark. Zool. 4 1953: 485–491 figs. 1–5.
- 4. Barnes, H. Considerazioni statistiche sulla distribuzione spaziale di alcuni organismi planctonici raccolti su un lungo percorso nel Golfo della Clyde. Mem. Ist., ital. Idrobiol. 7 1953: 109-127 fig. 1.

Barnes, H. see Rothschild, Lord.

- 5.—Barnett, R. C. Cell division inhibition of *Arbacia* and *Chaetopterus* eggs and its reversal by Krebs cycle intermediates and certain phosphate compounds. Biol. Bull. Woods Hole 104 1953: 263–274 figs. 1–10.
- 6.—Bataller, J. R. Equinodermos fosiles nuevos o poco conocidos de España. An. asoc. esp. Prog. Cienc. 6 1941: 60-73 figs.

Bennett, I. see Dakin, W. J.

7.—Bergounioux, F. M. Classe des Blastoïdes (Blastoïdea Say 1825) [in] Piveteau, J. Traité de Paléontologie 3 1953: 629-650 figs. 1-36.

- 8.—Bernasconi, I. Monografia de los Equinoideos Argentinos. An. Mus. Hist. nat. Montevideo (2) 6 1953: 1–58 pls. i–xxxii.
- 9.—Bernstein, M. H. & Mazia, D. The desoxyribonucleoprotein of sea urchin sperm. I. Isolation and analysis. II. Properties. Biochem. & Biophys. Acta 10 1953: 600–606 fig. 1; 11 1953: 59–68 figs. 1–4.
- 10.—Beyer, A. K. Zur Stratigraphie des obersten Gotlandiums in Mitteleuropa. Wiss. Z. Univ. Greifswald Math.-nat. R. 1 1951/1952: 35-67.

Björklund, U. see Borei, H. G.

- 11.—Blake, S. F. The Pleistocene fauna of Wailes Bluff and Langleys Bluff, Maryland. Smithson. misc. Coll. 121 12 1953: 1-32 fig. 1 pl. i. [Echinoderma at pp. 20, 23].
- 12.—Blum, H. F., Robinson, J. C. & Loos, G. M. The loci of action of ultraviolet and X-radiation and of photorecovery in the egg and sperm of the sea urchin Arbacia punctulata.

 J. gen. Physiol. 35 1951: 323-342 figs. 1-4 pl. i.

- 13.—Bock, K. J. & Schlieger, C. Über den Einfluss des Salzgehaltes im Meerwasser auf den Grundumsatz des Seesternes Asterias rubens L. Kieler Meeresforsch. 9 1953: 201-212 figs. 15-16.
- 14.—Borei, H. G. & Björklund, U. Effect of versine on sea urchin eggs. Exp. Cell Res. 5 1953: 216-219.
- 15.—Bowsher, A. L. A new Devonian crinoid from Western Maryland. Smithson. misc. Coll. 121 9 1953: 1–8 fig. 1 pl. i.

Bramlette, M. N. see Woodring. W. P.

Brewer, P. A. see Mazia, D.

- 16.—Broch, H. Ole Theodor Mortensen 22de februar 1868–3dje april 1952. K. norske vidensk. Selsk. Forh. 25 1953: 60–64 portrait.
- 17.—Buddenbrock, W. v. Vergleichende Physiologie Band 2. Nervenphysiologie. Basel (Verlag Birkhäuser) 1953: pp. 396 185 figs. [Echinoderma on pp. 105–132 figs. 51–70.]
- 18.—Cardin, A. & Meara, M. L. The lipids of sea-urchin semen. J. exp. Biol. 30 1953: 561-566.
- 19.—Čechovič, V. & Seneš, J. Torton okolia Modrého Kameňa. Geol. Sborn. Slovensk. Akad. 1 1950: 130-166 pls. xii-xvi [Summary in French at pp. 165-166.]
- 20.—Chace, F. A. Hubert Lyman Clark: Teacher and friend. January 9, 1870—July 31, 1947. Allan Hancock Pacif. Exped. 8 1948: ii—iv portrait.
- 21.—Chambers, E. L. & Mende, T. J. The adenosine triphosphate content of the unfertilized and fertilized eggs of Asterias forbesii and Strongylocentrous drobachiensis. Arch. Biochem. & Biophys. 42 1953: 45–56 fig. 1.
- 22.—Chambers, E. L. & Mende, T. J. Alterations of the inorganic phosphate and arginine phosphate content in sea urchin eggs following fertilization. Exp. Cell Res. 5 1953: 508-519 3 figs.

Chambers, E. L. see Mende, T. J.

23.—Cherbonnier, G. Recherches sur les Synaptes (Holothuries Apodes) de Roscoff. Arch. Zool. exp. gén. 90 1953 : 163-186.

- 24.—Cherbonnier, G. Note sur une nouvelle espèce de Synapte de l'Île Maurice: Patinapta vaughani n. sp. Bull. Mus. Hist. nat. Paris (2) 25 1953: 501-504 figs. a-m.
- 25.—Child, C. M. Redox indicator patterns in relation to echinoderm exogastrulation. 1. Oxidation patterns. Biol. Bull. Woods Hole 105 1953: 62-79 figs. 1-3.
- 26.—Child, C. M. Exogastrulation and differential cell dissociation by sodium azide in *Dendraster excentricus* and *Patiria miniata*. Physiol. Zoöl. 26 1953: 28–58 figs. 1–73.
- 27.—Chronic, J. [in] Newell, N. D., Chronic, J., and Roberts, T. G. Upper Paleozoic of Peru. Mem. geol. Soc. Amer. 58 1953: 1-230. [Echinoderma at pp. 164-165 pl. xxxv.]
- 28.—Clark, A. M. Notes on asteroids in the British Museum (Natural History). III Luidia. IV Tosia and Pentagonaster. Bull. Brit. Mus. (nat. Hist.) Zool. 1 1953: 379–412 figs. 1–15 pls. xxxix—xlvi.
- 29.—Clark, A. M. A revision of the genus *Ophionereis*. Proc. zool. Soc. Lond. 123 1953: 65-94 figs. 1-12 pls. i-iii.
- 30.—Clark, H. L. A report on the Echini of the warmer Eastern Pacific, based on the collections of the Velero III. Allan Hancock Pacif. Exped. 3 1948: 225-352 figs. 1-3 pls. xxxv-lxxi.
- 31.—Cleland, K. W. Heavy metals, fertilization and cleavage in the eggs of *Psammechinus miliaris*. Exp. Cell Res. 4 1953: 246-248.
- 32.—Comita, J. J. & Whiteley, A. H. The correlation of desoxyribonucleic acid synthesis and the rate of respiration in the sea urchin embryo. Biol. Bull. Woods Hole 105 1953: 412-419 figs. 1-2.
- 33.—Cooke, C. W. American Upper Cretaceous Echinoidea. Prof. Pap. U.S. geol. Surv. 254A 1953: 1–44 pls: i-xvi.
- 34.—Cornman, I. The effects of podophyllin on the maturation and cleavage of the starfish egg. Acta Un. int. cancr. 6 1949: 606–608 (reprinted as Contr. Bermuda biol. Sta. No. 154.)

- 35.—Cornman, I. Inhibition of seaurchin egg cleavage by a series of substituted carbamates. J. nat. Cancer Inst. 10 1950: 1123-1138 (reprinted as Contr. Bermuda biol, Sta. No. 158.)
- 36.—Cox, L. R. Cretaceous and Eocene fossils from the Gold Coast. Bull. geol. Surv. Gold Coast No. 17 1952: 1–68 pls. i-v.
- 37.—Creac'h, P. V. Sur la présence constante d'acide citrique dans le test calcaire des Invertébrés. C.R. Acad. Sci. Paris 237 1953: 944-946.

Crehuet, R. F. see Martin, O. R.

- 38.—Cuénot, L. Classe des Hétérostélés (Heterostelea Jackel 1918) [in] Piveteau, J. Traité de Paléontologie 3 1953: 599-606 figs. 1-12.
- 39.—Cuénot, L. Classe des Cystidés (Cystidea von Buch 1844) [in] Piveteau, J. Traité de Paléontologie 3 1953: 607-628 figs. 1-31.
- 40.—Dakin, W. J., Bennett, I. & Pope, E. Australian seashores. A guide for the beachlover, the naturalist, the shore fisherman, and the student. London (Angus & Robertson) 1953 pp. xii + 372 frontis. col. 99 pls. [Echinoderma at pp. 63, 310-335 pls. 79-93.]

Dan, K. see Mazia, D.

Danielli, J. F. see Lorch, I. J.

Danneberg, P. see Druckrey, H.

- 41.—Dartevelle, E. Echinides fossiles du Congo et de l'Angola. 2e partie. Description systématique des Echinides fossiles du Congo et de l'Angola. Ann. Mus. Congo belge 13 1953: xi, 1-240, 1 table pls. 1-19 figs. 1-56 pls. A-C, i-xix.
- 42.—Decourt, P., Anguéra, G. & Grenay, R. Effet de l'action narcobiotique sur le développement de l'œuf d'Oursin. C.R. Acad. Sci. Paris 236 1953: 2268-2271.
- 43.—Deflandre-Rigaud, M. Classe des Holothurides (Holothurioidea Zittel 1883) [in] Piveteau, J. Traité de Paléontologie 3 1953: 948-957 figs. 1-31.
- 44.—Dehm, R. Rhenechinus hopstätteri nov. gen. nov. sp., ein seeigel aus dem rheinischen Unter-Devon. Notizbl. hess. Landesamt. Bodenforsch. 81 1953; 88-95 figs. 1-2 pl. v.

- 45.—Deveze, L. Contribution a l'étude de la nutrition des stades larvaires planctoniques. II. Mise en evidence de l'utilisation possible du contenu bacterien des eaux par les plutei de Sphaerechinus granularis (Lamarck), Paracentrotus lividus Mortensen et d'Arbacia aequituber. (Blainville.) Rec. Trav. Sta. Mar. Endoume No. 8 1953: 55-59.
- 46.—Domantay, J. S. Littoral holothurians from Zamboanga and vicinity. Philipp. J. Sci. 82 1953: 109–131.
- 47.—Domantay, J. S. A brief summary of the Pacific and Atlantic Holothurioidea of the Allan Hancock Foundation of California. Philipp. J. Sci. 82 1953: 133–140.
- 48.—Druckrey, H., Danneberg, P., & Schmähl, D. Zur Pharmakologie zellteilungshemmender Gifte. (Versuche an Seeigeleiern.) Pubbl. Staz. zool. Napoli 24 1953; 247–266 figs. 1–5 pls. xii–xiii.
- 49.—Durham, J. W. Scutellaster and Anorthoscutum. J. Paleont. 27 1953: 147-149 fig. 1.
- 50.—Durham, J. W. Type species of Scutella. J. Paleont. 27 1953: 347–352 fig. 1 pl. 47.
- 51.—Durham, J. W. New name for Nipponaster Durham, 1952. J. Paleont. 27 1953: 756.
- 52.—Dyakonov, A. M. New species and subspecies of Asteroidea from the Far-East seas. Trav. Inst. Zool. Acad. Sci. U.S.S.R. 12 1952: 408-414.
- 53.—Eaton, T. H. Pedomorphosis; an approach to the chordate-echinoderm problem. Syst. Zool. 2 1953: 1–6 figs. 1–4.
- 54.—Endean, R. Queensland faunistic records. Part III. Echinodermata (excluding Crinoidea.) Pap. Dept. Zool. Univ. Qd. 1 1953: 51–60.
- 55.—Fell, H. B. Echinoderms from the Subantarctic Islands of New Zealand: Asteroidea, Ophiuroidea, and Echinoidea. Cape Expedition Series, Scientific Results of the New Zealand Subantarctic Expedition, 1941–45, No. 18. Dom. Mus. Rec. Zool., Wellington 2 1953: 73–111 figs. 1–6 pls. i–ii.

- 56.—Fell, H. B. The origin and migrations of Australasian echinoderm faunas since the Mesozoic. Trans. roy. Soc. N.Z. 81 1953: 245-255.
- 57.—Ficq, A. Incorporation in vitro de glycocolle- I_{-1} 4°C dans les occytes d'Astéries. Experientia 9 1953 : 377–379 figs. 1–2.
- 58.—Fontaine, A. The shallow-water echinoderms of Jamaica. Part I. The Starfishes (Class Asteroidea). Nat. Hist. Notes nat. Hist. Soc. Jamaica No. 59 1953: 179–184 8 figs.
- 59.—Fontaine, A. Shallow-water Echinoderms of Jamaica. Part II. The Brittle-stars (Class Ophiuroidea.) Nat. Hist. Notes nat. Hist. Soc. Jamaica No. 60 1953: 197–206 12 figs.
- 60.—Fontaine, A. The shallow-water echinoderms of Jamaica. Part III. The sea urchins (Class Echinoidea.) Nat. Hist. Notes nat. Hist. Soc. Jamaica No. 61 1953: 3-9 7 figs.
- 61.—Fontaine, A. The shallow-water echinoderms of Jamaica. Part IV. The sea-cucumbers (class Holothurioidea.) Nat. Hist. Notes nat. Hist. Soc. Jamaica Nos. 62–63 1953 : 29–33 7 figs.
- 62.—Geyer, O. F. Die Fauna der oolithischen Trümmerkalke des oberen Malm in Württemberg und ihre Beziehungen' zur korallogenen Fazies des Tithon. N. Jb. Geol. Paläont. B 3 1953: 130–140 [Echinoderma listed at p. 140.]
- 63.—Gigout, M. Études géologiques sur la Méséta Marocaine Occidentale (arrière-pays de Casablanca, Mazagan et Safi.) Trav. Inst. sci. Chérifien No. 3 1951: 1–508 pls. i–xxvii.
- 64.—Gill, W. D. Facies and fauna in the Bhadrar Beds of the Punjab Salt Range, Pakistan. J. Paleont. 27 1953; 824–844 figs. 1–3 pls. 88–91.

Grenay, R. see Decourt, P.

65.—Guislain, R. Recherches histochimiques sur les canaux de Cuyier de Holothuria impatiens Forskal. C.R. Soc. Biol. Paris 147 1953: 1254-1256.

Gustafson, T. see Lenicque, P.

66.—Harker, P. A new Edrioasteroid from the Carboniferous of Alberta. J. Paleont. 27 1953: 288–289 fig. 1.

- 67.—Harker, P. & Hutchinson, R. D. A new occurrence and redescription of Gogia prolifica Walcott. J. Paleont. 27 1953; 285–287 pl. xl figs. 1-5.
- 68.—Hassan, M. Y. [in] Donovan, D. T. De Danske Ekspeditioner til Østgrønland 1947–52. The Jurassic and Cretaceous stratigraphy and palaeontology of Traill Ø, East Greenland. Medd. Grønland 111 4 1953: 1–150 figs. 1–14 pls. i–xxv. [Echinoidea at pp. 128–129.]
- 69.—Hassan, M. Y. Tertiary faunas from Kap Brewster, East Greenland. Medd. Grønland 111 5 1953: 1-42 pls. A-E.
- 70.—Hatanaka, S. Notes on a Caudate Holothurian, Caudina similis (Augustin). Zool. Mag., Tokyo 51 1939; 347–353 figs. 1–6.
- . 71.—Hayasaka, I. On the echinoid called "Siraechinus kawaguchii." J. Fac. Sci. Hokkaido Univ. Geol. Min. 8 1953: 217-220 pl. xiv.
- 72.—Hayasaka, I. & Shibata, M. A new Tertiary species of *Echinarachnius* from Hokkaido. J. Fac. Sci. Hokkaido Univ. Geol. Min. 8 1952; 82–85 fig. 1.
- 73,—Haywood, C. The cleavage times of fertilized eggs of the sea urchin, Arbacia punctulata, at high pressures of nitrogen, helium, and nitrous oxide. J. cell. comp. Physiol. 41 1953: 335–343.
- 74.—Hirota, T. Physical analysis of the mechanism of cell division in sea urchin eggs. Publ. Seto mar. biol. Lab. 2 1953: 109–122 figs. 1–2 pl. iii.
- 75.—Holtedahl, O. Norges Geologi. Norges geol. Unders. 164 1953: 1–1117 78 figs.
- 76.—Hörstadius, S. The effect of lithium ions on centrifuged eggs of Paracentrotus lividus. Pubbl. Staz. zool. Napoli 24 1953 : 46-61 figs. 1-8.
- 77.—Hörstadius, S. & Runnström, J. Fertilization and membrane formation of sea urchin eggs aspirated in capillaries. Exp. Cell Res. 4 1953: 468–476 figs. 1–2.

Hörstadius, S. see Lenicque, P.

Hörstadius, S. see Lorch, I. J.

Hutchinson, R. D. see Harker, P.

- 78.—Lida, T. T. Changes of electric capacitance following artificial activation in sea urchin eggs. Zool. Mag., Tokyo 58 1949: 122–125. [English summary on p. 125.]
- 79.—Iida, T. T. Changes of electric capacitance following fertilization in sea urchin eggs.—Measurements over an extended frequency range. Zool. Mag., Tokyo 58 1949: 167–170. [English summary on p. 170.]
- 80.—Irwin, M. C. Sea urchins damage steel piling. Science 118 1953: 307.
- 81.—Ishida, G. Effect of KCN and other chemicals upon the action of the hatching enzyme in sea-urchin eggs. Zool. Mag., Tokyo 57 1947: 173–175. [English summary on p. 175.]
- 82.—Jacobson, F. W. & Millott, N. Phenolases and melanogenesis in the coelomic fluid of the echinoid Diadema antillarum Philippi. Proc. roy. soc. London 141B 1953: 231-247 figs. 1-6.
- 83.—Jeannet, A. Hessotiara zuberi sp. nov. Echinide nouveau du Jura soleurois. Vjschr. naturf. Ges. Zürich 98 1953: 176–177 figs. 1–5.
- 84.—Jensen, A. B. Some observations on the action of trypsin on sea urchin eggs. Exp. Cell Res. 4 1953: 60-68.
- 85.—Jensen, A. B. The effect of trypsin on the cross fertibility of sea urchin eggs. Exp. Cell Res. 5 1953: 325–328.
- **86.—Joysey,** K. A. A study of the type-species of the blastoid *Codaster* McCoy. Geol. Mag. **90** 1953: 208-218 figs. 1-5, 3 tables.

Kalckar, H. M. see Scarano, E.

- 87.—Kämpfe, L. & Kittel, R. Eine zoologische Fangreise in das Nordmeer, zugleich ein Beitrag zur fauna der Barents-See. Wiss. Z. Univ. Halle, Math.-nat. R. 2 1953: 463–469.
- 88.—Kavanau, J. L. Metabolism of free amino acids, peptides and proteins in early sea urchin development. J. exp. Zool. 122 1953: 285-337 figs. 1-11.
- 89.—Kennedy, G. Y. & Vevers, H. G. The biology of Asterias rubens L. V. A porphyrin pigment in the integument. J. Mar. biol. Ass. U.K. 32 1953: 235-247 figs. 1-2.

- 90.—Kennedy, G. Y. & Vevers, H. G. Protoporphyrin in the integument of Asterias rubens. Nature, London 171 1953: 31.
- 91.—Kerkut, G. A. The forces exerted by the tube feet of the starfish during locomotion. J. exp. Biol. 30 1953: 575-583 figs. 1-5.
- 92.—Kier, P. M. A new Lower Carboniferous echinoid from North America. Geol. Mag., London 90 1953: 65-69 figs. 1-4.

Kittel, R. see Kämpfe, L.

93.—Köster, E. Die Echiniden der tertiären Nahuel-Huapi-Serie von Nord-Patagonien. Bull. geol. Inst. Uniy. Upsala 34 1949–1953: 285–296 pl. i.

Koyasu, K. see Kuroda, C.

- 94.—Kriszat, G. Die Wirkung von Perjodat auf den Zustand der Plasmamembran des Seeigeleis. Exp. Cell Res. 5 1953: 420-426 1 fig.
- 95.—Kuhl, G. Die Wirkung von Colchiein auf die Dynamik der Furchung des Eies von Paummechinus miliaris Gmel. Verh. dtsch. zool. Ges. 1952 1953: 316–318.
- 96.—Kuhn-Schnyder, E. Vorgeschichtliche Tiere [in: Aargau: Natur und Erforschung.] Mitt. aargau. naturf. Ges. No. 24 1953: 207–223 figs. 1–11.

Küpper, K. see Papp, A.

- 97.—Kurian, C. V. A preliminary survey of the bottom fauna and bottom deposits of the Travancore coast within the 15-fathom line. Proc. nat. Inst. Sci. India 19 1953: 746-775 [Echinoderma at p. 756.]
- 98.—Kuroda, C. & Koyasu, K. The studies on the derivatives of Naphthoquinone. Part viii. The pigments from the sea urchins. iii. Proc. imp. Acad. Japan 20 1944: 23-25.
- 99.—Kuusi, T. Sur les effets des acides nucléiques et des protéines dans l'induction hétérogène. Arch. Biol., Paris 64 1953 : 189–226 figs. 1–8.
- 100.—Lafon, M. Récherches sur les sables côtiers de la Basse-Normandie et sur quelques conditions de leur peuplement zoologique. Ann. Inst. océanogr. Monaco N.S. 28 1953: 113-161.

- 6 [1953] 101 .- Laudon, L. R. & Severson, J. L.
- New Crinoid Fauna, Mississippian, Lodgepole Formation, Montana. Paleont. 27 1953: 505-536 figs. 1-2 pls. 51-55.
- 102.—Lawrence, L. J. The replacement of crinoid stems and gasteropods by cassiterite at Emmaville, New South Wales. J. roy. Soc. N.S.W. 86 1953: 119-122 pl, xii.
- 103.-Le Danois, E. La vie étrange des rivages marins. Paris (Horizons de France) 1953 pp. 192, 96 pls. (16 col.) text illust.
- 104.—Lenicque, P., Hörstadius, S. & Gustafson, T. Change of distribution of mitochondria in animal halves of sea urchin eggs by the action of micromeres. Exp. Cell Res. 5 1953: 400-403 3 figs.
- 105.—Lindahl, O. E. On a normally occurring reduction-division in somatic cells of the sea urchin embryo. Exp. Cell Res. 5 1953: 416-419 fig. 1.
- 106.—Lindahl, P. E. Somatic reduction division in the development of the sea urchin. Nature, London 171 1953: 437-438.
- 107.—Lindström, M. On the Lower Chasmops Beds in the Fågelsång District (Scania). Geol. Fören. Stockh. Förh. 75 1953: 125-148 figs. 1-6 pl. i.

Loos, G. M. see Blum, H. F.

- 108.—Lorch, I. J., Danielli, J. F. & Hörstadius, S. The effect of enucleation on the development of sea urchin eggs. I. Enucleation of one cell at the 2, 4 or 8 cell stage. II. Enucleation of animal or vegetal halves. Exp. Cell Res. 4 1953: 253-274 10 figs.
- 109.—Lowndes, A. G. The densities of some common Echinodermata from Plymouth. Ann. Mag. nat. Hist. (12) 6 1953 : 623-624.
- 110.—Lucas, G. Existence, dans l'appareil apical des Oursins, de deux sortes de plaques génitales. Plans de symétrie cristallographique. Hypothèses explicatives. C.R. Acad. Sci. Paris 237 1953: 405-407.
- 111.—Madsen, F. J. Asteroidea. Reports Swedish Deep-Sea Expedition 2 Zoology 1951: 73-92 figs. 1-2.
- 112,—Madsen, F. J. Ophiuroidea. Reports Swedish Deep-Sea Expedition 2 Zoology 1951: 107-117 figs. 1-2.

- 113.—Madsen, F. J. Holothurioidea. Reports Swedish Deep-Sea Expedition 2 Zoology 1953: 151-173 figs. 1-10.
- 114.—Maksimović, B. V. & Marković, B. Nouvelle contribution à la connaissance de l'Albien et du Cénomanien de la Serbie. Faune de l'Albien et du Cénomanien des environs de Kadina Luka et de Rajac (Serbie occidentale). Zborn. Radova geol. Inst. Beograd 5 1953: 183-208 figs. 1-4 pls. i-x. [Rec. Trav. Acad. Serbe Sci. 33 1953: 183-208 figs. 1-4 pls. i-x. Summary in French at pp. 208-210.]
- 115.-Maldonado-Koerdell, M. equinoides regulares del mesozoico de México. An. Escu. nat. Cienc. biol. Méx. 7 1953: 15–43 pls. i–ii.

Marković, B. see Maksimović, B. V.

116.-Marshak, A. & Marshak, C. Desoxyribonucleic acid in Arbacia eggs. Exp. Cell Res. 5 1953: 288-300 6 figs.

Marshak, C. see Marshak, A.

- 117.-Martin, O. R. & Crehuet, R. F. Apuntes para el estudio bionomico de la bahia de Santander. An. Assoc. esp. Progr. Cienc. 13 1948: 87-129 [See 63 in Zool. Rec. Vol. for 1948.]
- 118.-Marwick, J. Divisions and faunas of the Hokonui System (Triassic and Jurassic). Palaeont. Bull. N.Z. No. 21 1953 : 1-141 pls. i-xvii [Echinoderma at p. 47].
- 119.—Mazia, D., Brewer, P. A. & Alfert, M. The cytochemical staining and measurement of protein with mercuric bromphenol blue. Biol. Bull. Woods Hole 104 1953: 57-67 figs. 1-9.
- 120 .- Mazia, D. & Dan, K. isolation and biochemical characterization of the mitotic apparatus of dividing cells. Proc. nat. Acad. Sci., Wash. 38 1952: 826-838 figs, 1-2 pls. i-ii,

Mazia, D. see Bernstein, M. H.

Meara, M. L. see Cardin, A.

- 121.-Melendez, B. Los Carpoideos de España. An. Asoc. esp. Progr. Cienc. 17 1952 : 497-516 figs. 1-8 pls. i-ii.
- 122.—Mende, T. J. & Chambers, E. L. The occurrence of arginine phosphate in echinoderms. Arch. Biochem. and Biophys. 45 1953: 105-116.

Mende, T. J. see Chambers, E. L.

- 123.—Millott, N. Some preliminary observations on the young forms of the echinoid, *Diadema antillarum* Philippi. Bull. mar. Sci. Gulf and Caribbean 2 1953: 497-510 figs. 1-11.
- 124.—Millott, N. A remarkable association between Ophionereis reticulata (Say) and Harmothoe lunulata (Delle Chiaje). Bull. mar. Sci. Gulf and Caribbean 3 1953: 96–99 figs. 1–2.
- 125.—Millott, N. Colour pattern and the definition of the species, *Diadema* antillarum Philippi. Experientia 9 1953: 98-99 figs. 1-2.
- 126.—Millott, N. Observations on the skin pigment and amoebocytes, and the occurrence of phenolases in the coelomic fluid of *Holothuria forskali* delle Chiaje. J. Mar. biol. Ass. U.K. 31 1953: 529–539 fig. 1.
- 127.—Millott, N. Light emission and light perception in species of *Diadema*. Nature, London 171 1953: 973-974 figs. 1-3.
 - Millott, N. see Jacobson, F. W.
- 128.—Minato, M. Zur Orogenese und zum Vulkanismus im Jüngeren Palaeozoikum des Kitakami-Gebirges, Honshû, Japan. (Stratigraphische und Tektonische Untersuchungen des japanischen Palaeozoikums. Teil 6). J. Fac. Sci. Hokkaido Univ. Geol. Min. 7 1950: 277–302 pl. i.
- 129.—Minato, M. On some Palaeogene fossils in Hokkaido. J. geol. Soc. Japan 56 1950: 157-159 figs. 1-3.
- 130.—Minato, M. On the Lower Carboniferous fossils of the Kitakami Massif, Northeast Honsyû, Japan. J. Fac. Sci. Hokkaido Univ. Geol. Min. 7 1951: 355-372 pls. i-v.
- 131.—Minganti, S. The action of proteolytic enzymes on the vitelline membranes of *Psammechinus miliaris* eggs. Exp. Cell Res. 5 1953: 492–499 2 figs.
- 132.—Mitchison, J. M. The thickness of the sea urchin fertilization membrane. Exp. Cell Res. 5 1953: 536-538 1 fig.
- 133.—Mitchison, J. M. Microdissection experiments on sea urchin eggs at cleavage. J. exp. Biol. 30 1953: 515-524 pls. xiv-xv.

- 134.—Mitchison, J. M. & Swann, M. M. Measurements on sea-urchin eggs with an interference microscope. Quart. J. micr. Sci. 94 1953; 381-389 figs. 1-3.
 - Mitchison, J. M. see Swann, M. M.
- 135.—Monroy, A. A model for the cortical reaction of fertilization in the sea-urchin egg. Experientia 9 1953; 424-425.
- 136.—Monroy, A. & Oddo, A. M. Salubility changes in sea urchin eggs upon fertilization. J. gen. Physiol. 35 1951: 245-253 figs. 1-4.
- 137.—Moore, A. R. Alteration of developmental patterns in *Patiria miniata* by means of trypsin. J. exp. Zool. 123 1953: 561-569 pl. i.
- 138.—Moore, R. C. & Vokes, H. E. Lower Tertiary Crinoids from northwestern Oregon. Prof. Pap. U.S. Geol. Surv. 233—E 1953: 111-147 figs. 27-39 pls. 14-24.
- 139.—Morishita, A. Fossil Astriclypeus from Japan. Mem. Coll. Sci. Kyoto B 20 1952: 105-112 figs. 1-2 pl. xi.
- 140.—Morishita, A. On some Neogene Echinoids from Nagano Prefecture, Japan. Mem. Coll. Sci. Kyoto B 20 1953: 217–226 figs. 1-4, pl. i figs. 1-8.
- 141.—Morishita, A. Fossil species of the Palaeopneustidae from Japan. Trans. palaeont. Soc. Japan 9 1953: 27-29 pl. iii.
- 142.—Morishita, A. Neogene echinoids from Gifu Prefecture, Japan. Trans. palaeont. Soc. Japan 11 1953: 61-64 pl. vi.
- 143.—Mortensen, T. Echinoidea. Reports Swedish Deep-Sea Expedition 2 Zoology 1951: 47-48 fig. 1.
- 144.—Motomura, I. Corrective effect of x-dinitrophenol on the lithium larva of the sea urchin. Sci. Rep. Tôhoku Univ. (4) 18 1947: 81-84.
- 145.—Motomura, I. Secretion of sperm agglutinin in the fertilized eggs of the sea urchin. Exp. Cell Res. 5 1953: 187-190.
- 146.—Motomura, I. Secretion of the sperm agglutinin in the fertilized and denuded eggs of the sea urchin, Strongylocentrotus nudus. Sci. Rep. Tôhoku Univ. (4) 20 1953: 93-97.

- 147.—Müller, A. H. Bemerkungen zur Stratigraphie und Stratonomie der obersenonen Schreibkreide von Rügen. Teil H. Über die Bildungsgeschwindigkeit der Schreibkreide und die sich ergebenden Schlüsse. Geologie 2 1953: 25–34 figs. 1–5 pl. i.
- 148.—Müller, A. H. Die isolierten Skelettelemente der Asteroidea (Asterozoa) aus der obersenonen Schreibkreide von Rügen. Geologia Beihft 8 1953: 1-66 figs. 1-9 pls. i-xi.
- 149.—Murakami, S. A study on the dental and oral plates of the Ophiuroidea. Seibutu 2 1947: 97-104 figs. 1-2.
- 150.—Needham, J. Biochemistry and morphogenesis. Cambridge. 1950: i-xvi 1-787 figs. 1-328.
- 151.—Neefs, Y. Divers cas d'hermaphrodisme chez l'oursin Arbacia lixula L. Bull. biol. 87 1953: 461-468 figs. 1-6.
- 152.—Nieuwkoop, P. D. Concise catalogue of the Central Embryological Collection of the Hubrecht Laboratory, Utrecht. 1953 pp. 44. [Echinoid on p. 9.]

Oddo, A. M. see Monroy, A.

- 153.—Ohshima, H. Some rare abnormalities in Asteroids and Echinoids. Zool. Mag., Tokyo 51 1939: 158-162 figs. 1-2 pls. i-ii.
- 154.—Oka, T. B. Chromatoid fibrillar structure of the micropyle-canal of growing occytes of *Holothuria monacaria*. Zool. Mag., Tokyo 52 1940: 138-140 figs. 1-6.
- 155.—Palombi, A. & Santarelli, M. Gli animali commestibili dei mari d'Italia. Descrizione e nomi italiani, dialettali e stranieri dei pesci—tunicati—echinodermi—molluschi—crostacei. Milano (Ulrico Hoepli). 1953 pp. vi + 349 2 pls. col. 261 figs. [Echinoderma on pp. 213–216.]
- 156.—Pansini, R. Contributi sperimentali sur veleni antimitosica. II. L'influenza dell'etiluretano sulla morfogenesi delle uova di *Paracentrotus* lividus Lk. Arch. Sci. Biol., Napoli 35 1951: 339–359.
- 157.—Pansini, R. Contributi sperimentali sui veleni antimitosica. III. Influenza della Cloramina sulla morfogenesi delle uova di Paracentrolus lividus Lk. Bol. Soc. ital. Biol. sper. 28 1952: 1476–1477.

- 158.—Pantin, C. F. A. & Sawaya, P. Muscular action in *Holothuria grisea*. Bol. Fac. Filos. Sci. S. Paulo (Zool.) No. 18 1953 : 51-59 figs. 1-3.
- 159.—Papp, A. & Küpper, K. Holothurienreste aus dem Torton des Wiener Beckens. S.B. öst. Akad. Wiss. Abt. I 162 1953 ; 49-51 pl. i.
 - Parks, J. M. see Spreng, W. P.
- 160.—Patil, A. M. Study of the marine fauna of the Karwar coast. J. Bombay nat. Hist. Soc. 51 1953: 429-434 [Echinoderma on pp. 429-431].
- 161.—Pattee, M. Bibliography of Dr. H. L. Clark's Echinoderm papers. Allan Hancock Pacif. Exped. 8 1948: v-ix.
- 162.—Perlmann, P. Soluble antigens in sea urchin gametes and developmental stages. Exp. Cell Res. 5 1953: 394–399 7 figs.
- 163.—Pinar, N. & Roman, J. Echinides miocenes de la region de Tercan (Turquie). Rev. Fac. Sci. Univ. Istanbul 18B 1953: 173-182 fig.1 pl. i.
- 164.—Piveteau, J. Classe des Édrioastéroïdes (Edrioasteroidea Billings 1854-1858) [in] Piveteau, J. Traité de Paléontologie 3 1953 : 651-657 figs. 1-10.
 - Pope, E. see Dakin, W. T.
- 165.—Rees, C. B. The larvae of the Spatangidae. J. Mar. biol. Ass. U.K. 32 1953: 477-490 figs. 1-5.
- 166.—Reynolds, M. A. The Cainozoie succession of Maslin and Aldinga Bays, South Australia. Trans. roy. soc. S. Aust. 76 1953: 114-140 figs. 1-3.
- 167.—Richey, W. C. Vertebrate relatives. Proc. Pa Acad. Sci. 27 1953; 245-260 figs. 1-13.
- 168.—Robertson, J. D. Further studies on ionic regulation in marine invertebrates. J. exp. Biol. 30 1953: 277-296 fig. 1.
 - Robinson, J. C. see Blum, H. F.
- 169.—Roig, M. S. La geofisica en su relacion con la paleontologia. An. Acad. Habana 91 1953: 119-133.
- 170.—Roig, M. S. Nuevos Equinoideos fosiles de Cuba. An. Acad. Habana 91 1953: 135-176 pls. i-xii,

171.—Roig, M. S. Dos nuevos géneros de Equinoideos cubanos. Lambertona y Neopatagus, Mem. Soc. cubana Hist. nat. 21 1953: 257-262 pls. xxvii-xxix.

172.—Roig, M. S. Algunos Equinoideos fósiles Cubanos. Rev. Agric., Habana for 1953; pp. 53-67 pls. i-xxi [reprint].

173.—Roman, J. Sur les structures internes des Clypéastres. Bull. Soc. géol. Fr. 6 2 for 1952 1953 : 403-416 fig. 1.

Roman, J. see Pinar, N.

174.—Rösler, A. Agelacrinites taunicus n. sp. (Edrioaster.) aus dem Unterdevon von Mensfelden (Blatt Limburg, Lahn). Notizbl. hess. Landesamt. Bodenforsch. 81 1953: 69-74 pl. iii.

175.—Rothschild, Lord. The fertilization reaction in the sea-urchin. The induction of polyspermy by nicotine. J. exp. Biol. 30 1953: 57-67 figs. 1-4 pl. v.

176.—Rothschild, Lord & Barnes, H. The inorganic constituents of the sea urchin egg. J. exp. Biol. 30 1953; 531-541

177.—Rudenberg, F. H. N. The role of the jelly coat in the uptake of calcium by eggs of *Arbacia punctulata* before and after fertilization. Exp. Cell Res. 4 1953: 116-126 figs. 1-3.

Runnström, J. see Hörstadius, S.

178.—Russo, G. La basi esoniche e puriniche libere nel testicolo di *Paracen*trotus lividis, durante il ciclo funzionale. Bol. Soc. ital. Biol. sper. 28 1952: 757-758.

Sanchez Roig, M. see Roig, M. S.

Santarelli, M. see Palombi, A.

Sawaya, P. see Pantin, C. F. A.

179.—Scarano, E. & Kalekar, H. M. Nucleic acid synthesis in developing sea urchin embryos. Pubbl. Staz. zool. Napoli 24 1953: 188–196.

Schlieger, C. see Bock, K. J.

Schmähl, D. see Druckrey, H.

180.—Seilacher, A. Studien zur Palichnologie II. Die fossilen Ruhespuren (Cubichnia). Neues Jb. Min. Geol. Paläont. 93 1953: 87-124 figs. 1-5 pls. vii-xiii. Seneš, J. see Čechović, V.

181.—Severin, K. A starfish rights itself. Nat. Hist., N.Y. **62** 1953: 276-277 figs. 1-7.

Severson, R. see Laudon, L. R.

Shibata, M. see Hayasaka, I.

182.—Shikama, T. A handbook of illustrated fossils from Japanese and their adjacent territories. Kyôto, 1952: 1-376 figs. 1-74 pls. i-lvii.

183.—Shrock, R. R. & Twenhofel, W. H. Principles of Invertebrate Paleontology. 2nd ed. 1953 [Echinoderma at pp. 642-735 figs. (14) 1-57.]

184.—Sieverts-Doreck, H. Über Austinocrinus im norddeutschen Senon. Mit einem Beitrag zur Gliederung, Stammesgeschichte und Verbreitung der Gattung. Mitt. geol. (St.) Inst. Hamb. 22 1953: 102-118 figs. 1-6 pls. 17-18.

185.—Sieverts-Doreck, H. Zur Verbreitung känozoischer Ophiuren. N. Jb. Geol. Paläont. B 6 1953: 275–286.

186.—Sieverts-Doreck, H. Über einige inadunate Crinoiden aus dem rheinischen Devon. Notizbl. hess. Landesamt. Bodenforsch. 81 1953: 75-87 figs. 1-8 pl. iv.

187.—Sikošek, B. Stratigraphie und Tectonik des gebeites zwischen Mali Izoor—Minćevo—Kadibogaz in Ostserbien. Zborn. Radova geol. Inst. Beograd 4 1952: 146–167 (Summary in German at p. 166) pls. i–iii [Rec. Trav. Acad. Serbe Sci. 23 1952: 146–167].

188.—Spreng, W. P. & Parks, J. M. Evolution in basal plates of monocyclic camerate crinoids. J. Paleont. 27 1953: 585-595 figs. 1-8.

189.—Strimple, H. L. A new species of Archaeocrinus from Oklahoma. J. Paleont. 27 1953: 604-606 figs. 1-7.

190.—Strimple, H. L. A new carpoid from Oklahoma. J. Wash. Acad. Sci. 43 1953: 105–106 figs. 1–2.

191.—Strimple, H. L. A new species of Carinocrinus from Oklahoma. J. Wash. Acad. Sci. 43 1953: 201-203 figs. 1-2.

192.—Sugiyama, M. Studies on the artificial parthenogenesis of the sea urchin eggs. VIII. Zool. Mag., Tokyo 58 1949: 137-138 fig. 1.

- 193.—Sugiyama, M. Studies on the artificial parthenogenesis of the sea urchin eggs. IX. The lowest effective concentration of fatty acids for membrane formation. Zool. Mag., Tokyo 58 1949: 156–157. [English summary on p. 157.]
- 194.—Sugiyama, M. Studies on the artificial parthenogenesis of the sea urchin eggs. X. Effect of Body Fluid. Zool. Mag., Tokyo. 58 1949: 159–160. [English summary on p. 160.]
- 195.—Sugiyama, M. Studies on the artificial parthenogenesis in the sea urchin eggs. XI. Effects of salt ions on the membrane formation. Zool. Mag., Tokyo 58 1949: 160–162. [English summary on p. 162.]
- 196.—Sugiyama, M. Studies on the artificial parthenogenesis in the sea urchin eggs. XII. Effect of Ca-defect. Zool. Mag., Tokyo 58 1949: 183–184. [English summary on p. 184.]
- 197.—Sugiyama, M. Studies on the artificial parthenogenesis in the sea urchin eggs. XIII. Membrane formation by copper sulfate and corrosive sublimate. Zool. Mag., Tokyo 58 1949: 223–224 fig. [English summary on p. 224.]
- 198.—Sugiyama, M. Polyspermy in the sea-urchin eggs induced by CuSO₄. Zool. Mag., Tokyo 59 1950: 11-12. [English summary at p. 12.]
- 199.—Sugiyama, M. Studies on the artificial parthenogenesis in the sea urchin eggs. XIV. The method to sensitize the eggs for membrane formation by distilled water or urea solution. Zool. Mag., Tokyo 59 1950: 78–79. [English summary on p. 79.]
- 200.—Sugiyama, M. Studies on the artificial parthenogenesis in the sea urchin eggs. XV. Membrane formation by formalin. Zool. Mag., Tokyo 59 1950: 100-101.
- 201.—Sugiyama, M. Studies on the artificial parthenogenesis in the sea urchin eggs. XVI. Structure of the fertilization membrane and its formation. Zool. Mag., Tokyo 59 1950: 102–103 figs. [English summary on p. 103.]

- 202.—Sugiyama, M. Studies on the artificial parthenogenesis in the sea urchin eggs. XVII. The conditions to induce the membrane formation by means of saponin. Zool. Mag., Tokyo 59 1950: 132–134. [English summary on p. 134.]
- 203.—Sugiyama, M. Studies on the artificial parthenogenesis in the sea urchin eggs. XVIII. Membrane formation by means of urea albumen solution. Zool. Mag., Tokyo 59 1950: 135–136. [English summary on p. 136.]
- 204.—Sugiyama, M. Studies on the artificial parthenogenesis in the sea urchin eggs. XIX. Membrane formation by means of hypertonic solution. Zool. Mag., Tokyo 59 1950: 158-159. [English summary on p. 159.]
- 205.—Sugiyama, M. Studies on the artificial parthenogenesis in the sea urchin eggs. XX. Membrane formation by Na-choleinate. Zool. Mag., Tokyo 59 1950: 160-162. [English summary at p. 162.]
- 206.—Sugiyama, M. Studies on the artificial parthenogenesis in the sea urchin eggs. XXI. Membrane formation by means of Alcohols. Zool. Mag., Tokyo 60 1951: 124—126. [English summary on p. 126.]
- 207.—Sugiyama, M. Physiological analysis of the cortical response of the sea urchin egg to stimulating reagents. I. Response to sodium choleinate and wasp-venom. Biol. Bull. Woods Hole 104 1953: 210–215 figs. 1–2.
- 208.—Sugiyama, M. Physiological analysis of the cortical response of the sea urchin egg to stimulating reagents. II. The propagating or non-propagating nature of the cortical changes induced by various reagents. Biol. Bull. Woods Hole 104 1953: 216–223 figs. 1–4.
- 209.—Sun, Y. C. & Szetu, S. S. The stratigraphical and biological position of the species "Camarocrinus asiaticus". Bull. geol. Soc. China 27 1947: 243–252 figs. 1–5 pl. i.
- 210.—Swan, E. F. The Strongylocentrotidae (Echinoidea) of the Northeast Pacific. Evolution 7 1953: 269-273.
- 211.—Swann, M. M. Structural factors in mitosis. Abstr. Diss. Univ. Camb. 1950-51 1953: 35-38.

- 212.—Swann, M. M. The mechanism of cell-division. A study with carbon monxide on the sea-urchin egg. Quart. J. micr. Sci. 94 1953: 369-379 figs. 1-7.
- 213.—Swann, M. M. & Mitchison, J. M. Cleavage of sea-urchin eggs in colchicine. J. exp. Biol. 30 1953: 506– 514 figs. 1–4 pls. xii–xiii.

Swann, M. M. see Mitchison, J. M.

Szetu, S. S. see Sun, Y. C.

- 214.—Szörényi, E. Moizäne Echinoiden aus den westlichen teilen der Ukraine. Geol. hung. Ser. Palaeont. 23 1953: 1-104 pls. i-viii. [German translation pp. 49-102.]
- 215.—Tartarin, A. Observations sur les mutilations, la regeneration, les neoformations, et l'anatomie de Coscinasterias tenuispina Lamarck. Rec. Trav. St. mar. Endoume No. 10 1953: 1-107 figs. 1-57.

Termier, G. see Termier, H.

- 216.—Termier, H. & Termier, G. Généralités sur les Échinodermes [in] Piveteau, J. Traité de Paléontologie 3 1953:587-598 figs. 1-7.
- 217.—Termier, H. & Termier, G. Classe des Échinides (Echinoidea Zittel 1883) [in] Piveteau, J. Traité de Paléontologie 3 1953: 857-947 figs. 1-250 pls. i-v.
- 218.—Tokioka, T. Invertebrate fauna of the intertidal zone of the Tokara Islands. V. Echinodermata. Publ. Seto mar. biol. Sta. 3 1953: 144-148 figs. 1-3 pl. vii.
- 219.—Tortonese, E. Nota preliminare intorno ad alcuni Echinodermi nuovi per il golfo di Genova e le zone vicine. Atti Accad. Ligure 7 1951: 268-272.
- 220.—Tortonese, E. Gli Echinodermi del Mar Ligure e delle zone vicine. Atti Accad. Ligure 8 1952: 163-242 figs. 1-10 pl. i.
- 221.—Tortonese, E. On Lytechinus (Echinoidea) from the West Coast of South America. Ann. Mag. nat. Hist. (12) 6 1953: 479-480.
 - Twenhofel, W. H. see Shrock, R. R.

- 222.—Tyler, A. Prolongation of life-span of sea-urchin spermatozoa, and improvement of the fertilization-reaction, by treatment of spermatozoa and eggs with metal-chelating agents (amino acids, Versene, DEDTC, oxine, cupron). Biol. Bull. Woods Hole 104 1953: 224–239 figs. 1–8.
- 223.—Ubaghs, G. Notes sur Lichenoides priscus Barrande, eocrinoide du Cambrien Moyen de la Tchécoslovaquie. Bull. Inst. Sci. nat. Belg. 29 34 1953: 1-24 figs. 1-12.
- 224.—Ubaghs, G. Classe des Crinoides (Crinoidea Miller 1821) [in] Piveteau, J. Traité de Paléontologie 3 1953: 658-773 figs. 1-166.
- 225.—Ubaghs, G. Classe des Stelléroides (Stelleroidea) [in] Pivetean, J. Traité de Paléontologie 3 1953: 774–842 figs. 1–64.
- 226.—Ubaghs, G. Classe des Ophiocistioides (Ophiocistioidea) [in] Piveteau, J. Traité de Paléontologie 3 1953: 843–856 figs. 1–8.
- 227.—Ubisch, L. von. Über seeigelmerogone. Experientia 9 1953: 294–295 figs. 1–4.
- 228.—Urbani, E. Osservazioni spettrofotometriche sull'oocite in crescita di "Antedon rosacea" Lamarck. R.C. Accad. Lincei (8) 15 1953: 308–315 figs. 1–8.
- 229.—Valensi, L. Microfossiles des silex du Jurassique moyen. Remarques petrographiques. Mem. Soc. géol. France No. 68 1953: 1-100 figs. 1-4 pls. i-xvi.
- 230.—Vannini, E. Osservazioni sull'accrescimento del nucleolo negli ovociti di Antedon rosacea. Pubbl. Staz. zool. Napoli 24 1953: 292–331 figs. 1–9.
- 231.—Vasseur, E. & Wicklund, E. Correlation of some physiological properties of the jelly coat solution of sea-urchin egg. Ark. zool. 4 1953: 363–366.

Vevers, H. G. see Kennedy, G. Y.

232.—Vincent, W. S. The isolation and chemical properties of the nucleoli of starfish oocytes. Proc. nat. Acad. Sci., Wash. 38 1952: 139-145.

233.—Vinogradov, A. P. The elementary chemical composition of marine organisms. Mem. Sears Fdn. Mar. Res. 2 1953: 1-647. [Echinoderma on pp. 243-270 tables 142-161.]

Vokes, H. E. see Moore, R. C.

234.—Vyalov, O. S. [The problem of the classification of Crinoid columnals.] C.R. Acad. Sci. URSS n.s. 89 1953: 1087-1090.

235.—Walier, J. C. Paleontology of Rustler Formation, Culberson County, Texas. J. Paleont. 27 1953: 679-702 pls. lxx-lxxiii.

236.—Weel, P. B. van. Mechanical properties of the resting holothurian muscle. Proc. Hawaii, Acad. Sci. No. 28 1953: 6-7.

237.—Wells, P. H. Photoreactivation in gametes of the sea urchin Strongylocentrous purpuratus. Stanf. Univ. Bull. 27 1953: 164-165.

238.—Westblad, E. New Turbellaria parasites in Echinoderms. Ark. Zool. 5 1953: 269-288 figs. 1-9 pls. i-iii.

Whiteley, A. H. see Comita, J. J.

Wicklund, E. see Vasseur, E.

239.—Wilcox, N. R. Zonal variation in selected morphological features of *Echinocorys scutata* Leske. Geol. Mag., London 90 1953: 83–96 figs. 1–6.

240.—Wind, J. Kridtaflejringer i Jylland. Flora og Fauna 59 1953: 73-84.

241.—Woodring, W. P. & Bramlette, M. N. Geology and paleontology of the Santa Maria district, California. Prof. Pap. U.S. Geol. Surv. 222 1950: 1-142 figs. 1-8 pls. 1-23.

242.—Yakovlev, N. N. [Self-adjustment and development of forms in sea lilies]. C.R. Acad. Sci. URSS n.s. 86 1952: 827-828.

243.—Yanagimachi, R. A note on the so-called "Boring movement" of sea urchin spermatozoa. Zool. Mag., Tokyo 62 1953: 220-224 figs. 1-3. [English summary at p. 224.]

244.—Zeuthen, E. Biochemistry and metabolism of cleavage in the sea urchin egg, as resolved into its mitotic steps. Colloq. Union int. Sci. Biol. No. 12 1953: 31-58 7 figs.

245.—Zoeke, M.-E. Heterosalenia alloiteaui nov. sp. du Jurassique moyen du Liban Nord et un cas croissance excessive de plaques arrêtant le développement des zones ambulacraires. Bull. Soc. géol. Fr. 6 2 for 1952 1953: 249-252 text-figs. 1–2 pl. xib.

H.—SUBJECT INDEX

GENERAL LITERATURE AND HISTORY

Text-books; General Works.—Ber-Goundux 7; Cuenot 38-39; De-Flaxdre-Ricaud 43; Le Danois 103; Needham 150; Piveteau 164; Shikama 182 pp. 213-219 figs. 19-23; Serock & Twenhopel 183; Termier & Termier 216-217; Ubaghs 224-226.

Biography: Obituary.—H. L. Clark, CHACE 20; Mortensen, BROCH 16.

Bibliography.—H. L. Clark, PATTEE 161.

Nomenclature.—Classification of fossil crinoid columnals, Vyalov 234.

Museums: Collections. — Hubrecht Laboratory, Utrecht, NIEUWKOOP 152.

Vernacular names.—Italian Echinoidea, Palombi & Santarelli 155.

Geophysical methods of dating fossil Echinoderma.—Roug 169.

STRUCTURE

Skeletal Anatomy.—Terminology of Asteroid plates, Müller 143; dental and oral plates, Ophiuroidea, Mura-Kami 149; internal structures in Clypeaster, Roman 173; crystallographic orientation of genital plates in Echinoidea, Lucas 110; Yakovlky 242.

Circulatory Systems.—Guislain 65; Robertson 168.

Body Cavity.—Pyloric caeca, Asterias forbesi, Anderson 2.

Nervous System.—Buddenbrock 17.

Reproductive System.—OKA 154.

Special Structures.—Iridophores in Diadema antillarum, MILLOTT 127.

Teratology.—Ohshima 153; Zoeke 245.

Laboratory Technique. — Tartarin 215.

PHYSIOLOGY

Nutrition .- DEVEZE 45.

Density.—Lowndes 109.

Autotomy and Regeneration. — Linckia, [ANON.] 1; Coscinasterias, Tartarin 215.

Chemical Constituents.—Bernstein & Mazia 9; semen of Echinus esculentus, Cardin & Meara 18; Chambers & Mende 21; citric acid in tests of Paracentrotus lividus and Sphaerechinus granularis, Creac'h 37; Kennedy & Vevers 89-90; echinoid radioles, Kuroda & Koyasu 98; Mende & Chambers 122; melanin and tyrosinase in Holothuria, Millott 126; eggs of Paracentrotus lividus, Rothschild & Barnes 176; nucleolus from oocytes of Asterias rubens, Vincent 232; Vinogradov 233.

Coloration: Colour changes.—Diadema, MILLOTT 123, 125.

Muscular Activity.—Pantin & Sawaya 158.

Locomotion; movement. — Asterias rubens, Kerkut 91.

Luminosity.—Iridophores in Diadema antillarum, MILLOTT 127.

Experimental.—Bäckström 3: Bar-NETT 5; BLUM, ROBINSON & LOOS 12; BOCK & SCHLIEGER 13; BOREI & Björklund 14; Buddenbrock 17; Cardin & Meara 18; Chambers & Mende 22; CHILD 25-26; CLELLAND 31; Comita & Whiteley 32; Decourt, Anguéra & Greenay 42; Duckbey, DANNEBERG & SCHMÄHL 48; FICQ 57; HAYWOOD 73; HIROTO 74; HÖRSTADIUS 76; HÖRSTADIUS & RUNNSTRÖM 77; IIDA 78-79; ISHIDA 81; JACOBSON & MILLOTT 82; JENSEN 84-85; KAVANAU 88; KRISZAT 94; KUHL 95; KUUSI 99; Lenicque, Hörstadius & Gustafson 104; Lorch, Danielli & Hörstadius 108; Marshak & Marshak 116; Mazia & Dan 120; Minganti 131; Mitchison 132; MITCHISON & SWANN 134; MONROY 135; Monroy & Oddo 136; Motomura 145-146; PANSINI 156-157; PANTIN & SAWAYA 158; PERLMANN 162; ROTHS-CHILD 175: ROTHSCHILD & BARNES 176: RUDENBERG 177; RUSSO 178; SCARANO & Kalckar 179; Sugiyama 192-208; Swann 210, 212; Tyler 222; Urbani 228; Vannini 230; Vasseur & Wicklund 231; Weel 236; Wells 237; Zeuthen 244.

Laboratory Technique. — Mazia, Brewer & Alfert 119.

REPRODUCTION AND SEX

Hybridization.—Swann 210; Ubisch 227

Parthenogenesis.—Sugiyama 192-206.

Hermaphrodism .- NEEFS 151.

DEVELOPMENT

Fertilization.—Yanagımachı 243.

Larval Stages.—Paracentrotus lividus, Strongylocentrotus droebachiensis, Lin-Dahl. 105 & 106; Brissopsis lyrifera, Echinocardium cordatum, E. flavescens, Spatangus purpureus, REES 165 figs.

Post-Larval Stages.—Codaster, Joysey 86; colour changes in Diadema antillarum, Millott 125.

Experimental Embryology.—Barnett 5; Blum, Robinson & Loos 12; Child 25; Cleiand 31; Corman 34-35; Druckrey, Danneberg & Schmähl 48; Hörstadius & Runström 77; Jensen 34; Kuhl 95; Lorch, Danielli & Hörstadius 108; Mitchison 133; Mitchison & Swann 134; Monroy 135; Moore 137; Motomura 144-146; Rottischild 175; Scarano & Kalckar 179; Swann 211-212; Swann & Mitchison 213; Yanagimachi 243; Zeuthen 244.

EVOLUTION AND GENETICS

General Works: Theories.—EATON 53.

Phylogeny.—RICHEY 167; Austinocrinus, SIEVERTS-DORECK 184; basal plates in monocyclic camerate crinoids, SPRENG & PARKS 188.

Variation .- Echinocorys, WILCOX 239.

ECOLOGY AND HABITS

Communities.-LAFON 100.

Habitat.—SWANN 210; Echinocorys, Wilcox 239.

Distribution.—Spp. with a bathymetrical distribution to 3000 metres or below, Madsen 111-113, Mortensen 143.

Parasites and Commensals.—Ophionereis reticulata and polychaete Harmothoe lunulata, MILLOTT 124; Turbellaria in Echinoderma, Westblad 238.

Boring Habits.—Strongylocentrotus purpuratus boring into steel piles, IRWIN 80.

Locomotion.—Kerkut 91; Severin 181.

Economics.—Italian Echinoidea as food for man, Palombi & Santabelli 155.

Geological Aspect.—Echinoderma as a guide to rate of deposition of Rügen Chalk, MÜLLER 147.

DISTRIBUTION

GEOGRAPHICAL

General.—Relation between recent and Tertiary faunas of Australasia, Fell 56.

Arctic.-DYAKONOV 52.

North Atlantic.—Barnes 4; Chebbonnier 23; A. M. Clark 28; Kampfe & Kittel 87; Lafon 100; Le Danois 103; Madden 111-113; Martin & Crehuet 117; Mortensen 143; Rees 165.

Mediterranean and Black Sea.—A. M. CLARK 28; LE DANOIS 103; PALOMBI & SANTARELLI 155; TORTONESE 219-220.

North Pacific.—A. M. CLARK 28-29; H. L. CLARK 30; DOMANTAY 47; HATANAKA 70; HAYASAKA 71; IRWIN 80; MORISHITA 139.

Tropical Atlantic.—A. M. CLARK 29; DARTEVELLE 41; DOMANTAY 46; FON-TAINE 58-61; MADSEN 111-113; MIL-LOTT 123-124.

Indo-Pacific. — CHERBONNIER 24; A. M. CLARK 28-29; H. L. CLARK 30; DAKIN, BENNETT & POPE 40; DOMAN-TAY 46-47; ENDEAN 54; FELL 56; KUELAN 97; PATIL 160; TOKIOKA 218; TORTONESE 221.

Polynesian Seas.—A. M. CLARK 29; DOMANTAY 47.

Eastern Pacific.—A. M. CLARK 29; DOMANTAY 47.

South Temperate and Subantarctic.— BEENASCONI 8; A. M. CLARK 28-29; DAKIN, BENNETT & POPE 40; ENDEAN 54; FELL 55-56.

GEOLOGICAL

Pleistocene.—U.S.A., BLAKE 11; Angola, Belgian Congo, Dartevelle 41; Australia, Lawrence 102; Japan, Shikama 182; Sievers-Doreck 185.

Tertiary.—Malta, Durham 50; Patagonia, Köster 93; Australia, Reynolds

Neogene.—Durham 51; Greenland, Hassan 69; Japan, Morishita 142; Japan, Shikama 182.

Pliocene.—U.S.A., DURHAM 49; Australia, Fell 56; Formosa, Hayasaka 71; Japan, Morishita 140–141; Australia, Reynolds 166; Sieverts-Doreck 185; U.S.A., Woodring & Bramlette 241.

Miocene.—Czechoslovakia, Čechovič & Seneš 19; Angola, Belgian Congo, Daetevelle 41; Dueham 50; Australasia, Fell 56; Japan, Havasuka & Shibata 72; Japan, Morishita 139-140; Austria, Papp & Küpper 159; Turkey, Pinar & Roman 163; Cuba, Roig 170, 172; Sleverts-Doreck 185; Ukraine, Szörényi 214; U.S.A., Woodring & Bramlette 241.

Palaeogene.—Japan, Minato 129; Japan, Shikama 182.

Oligocene.—Australasia, Fell 56; U.S.A., Moore & Vokes 138; Cuba, Roig 170-172; Sieverts-Doreck 185.

Eccene.—Gold Coast, Cox 36; Angola, Belgian Congo, Darrevelle 41; Australasia, Fell 56; Pakistan, Gill 64; Cuba, Roic 170-172; Sieveetts-Dorbek 185; Denmark, Wind 240.

Cretaceous. — Spain, Bataller 6; America, Cooke 33; Angola, Belgian Congo, Dartevelle 41; Australasia, Fell 56; N. Africa, Gigout 63; Greenland, Hassan 68; Yugoslavia, Maksimović & Marković 114; Mexico, Maldonado-Koerdeill 115; Rügen, Müller 147–148; Cuba, Roic 170; Sieverts-Doreck 184; England, Wilcox 239; Denmark, Wind 240.

Jurassic.—Spain, Bataller 6; Germany, Gever 62; Switzerland, Jeannet 83; Switzerland, Kuhn-Schnyder 96; Mexico, Maldonado-Koerdell 115;

Seilacher 180; Japan, Shikama 182; Yugoslavia, Sikošek 187; France, Valensi 229; N. Africa, Zoeke 245.

Permian.—Peru, Chronic 27; Australia, Lawrence 102; Texas, Walter 235.

Trias. — Switzerland, Kuhn-Schnyder 96; New Zealand, Marwick 118.

Carboniferous.—Peru, Chronic 27; Canada, Harker 66; England, Joysey 86; Canada, Kier 92; U.S.A., LAUDON & Severson 101; Japan, Minato 128, 130; U.S.A., Strimple 191.

Devonian.—U.S.A., BOWSHER 15; Germany, DEHM 44; Germany, RÖSSLER 174; Germany, SIEVERTS-DORECK 186.

Silurian.—Europe, Beyer 10; U.S.A., Bowsher 15; China, Sun & Szetu 209.

Ordovician. — Norway, Holtedahl 75; Sweden, Lindström 107; U.S.A., Strimple 189-190.

Cambrian. — Canada, HARKER & HUTCHINSON 67; Spain, MELENDEZ 121; Czechoslovakia, UBAGHS 223.

III.—SYSTEMATIC INDEX

Full references are given only for new genera or species, and the volume number is printed in "small bold Clarendon." In other references the numeral in "large bold Clarendon" refers to the list of titles where the full reference will be found.

HOLOTHURIOIDEA

Actinopyga echinites p. 120, miliaris p. 120, serratidens p. 121, Domantay 46.—A. agassizii, Fontaine 61 p. 30 fig.—A. mauritiana, Tokioka 218 p. 147 pl. vii figs. 5-7.

Astichopus multifidus, Fontaine 61 p. 31.

Benthodytes typica, Madsen 113 p. 160 fig. 8.

Caudina similis, HATANAKA 70 p. 347 figs. 1-6.

Chiridota peloria, rotifera, Fontaine 61 p. 32.

†Chiridotites vetustus p. 70 pl. xi figs. 9, 12, C. sp. p. 70 pl. xi fig. 13, VALENSI 229.

Cucumaria cucumer p. 232, planci p. 229, syracusana p. 230 fig. 10, tergestina p. 231 pl. i fig. 10, C. sp. p. 230 fig. 9, TORTONESE 220.

†Deimatidae spicules, PAPP & KÜP-PER 159 p. 51 pl. i figs. 1-4, 5 ?

Euapta lappa, Fontaine 61 pp. 29, 32 figs.

Euphronides kerhervei, MADSEN 113 p. 161 fig. 9.

Holothuria saecularis var. forskali var. n. Philippine Is., Domantay Philipp. J. Sci. 82 pp. 118-119.—H. impatiens, parvula, Fontaine 61 p. 30 figs.—H. forskali, coelomic fluid, Millott 126.—H. forskali p. 227, helleri p. 227, impatiens p. 224, polii p. 226, sanctori p. 225, tubulosa p. 225, Tortonese 220.—H. forskali, impatiens, polii, tubulosa, Turbellaria parasitic in, Westblad 238.

Holothuria (Holothuria) atra p. 147 pl. vii figs. 1-2, vagabunda p. 146 pl. vii figs. 3-4, ТОКІОКА 218.

Leptosynapta cruenta sp. n, N. France, CHERBONNIER Arch. Zool. exp. gén, 90 pp. 175-179 pl. iv figs. a-j; galliennei p. 168 pl. ii, inhaerens p. 173 pl. iii, minuta p. 165 pl. i, ID.—L. dolabrifera, DAKIN, BENNETT & POPE 40 p. 334 pl. xciii figs. 4-5.

Mesothuria candelabri, Madsen 113 p. 153.—M. verrilli, Tortonese 220 p. 228.

Microthele nobilis, Domantay 46 p. 122.

†Molpadiidae spicules, PAPP & KÜP-PER 159 p. 51 pl. i figs. 6-8.

†Myriotrochites cf. costiferus, VALENSI 229 p. 70,

Oestergrenia digitata, Tortonese 220 p. 233.

Opheodesoma spectabilis var. puertogalerae, Domentay 46 p. 127.

Patinapta vaughani sp. n. Mauritius, Cherbonnier Bull. Mus. Hist. nat. Paris (2) 25 pp. 501-504 figs, a-m. Peniagone nybelini sp. n. Tropical Atlantic, Madsen Rep. Swedish Deep-Sea Exped. 2 pp. 157-158 fig. 5; lugubris p. 153 figs. 2-3, P. sp. (? lugubris) p. 155, porcellus p. 155 fig. 4, P. (or Scotoanassa) sp. p. 159 fig. 7, In

Phyllophorus seguroensis, Fontaine 61 p. 31.—P. fragilis, Tokioka 218 p. 147 fig. 3, pl. vii fig. 15.

Polycheira rufescens, Tokioka 218 p. 147 pl. vii figs. 16-17.

Protankyra brychia, Madsen 113 p. 151 fig. 1.

Pseudocucumis sp., Tokioka 218 p. 147 pl. vii figs. 12-14.

Psolidium nigrescens, DAKIN, BENNETT & POPE 40 p. 335 pl. xciii figs. 2-3.

Psychropotes aff. minutus, Madsen 113 p. 163 fig. 10.

Scotoanassa translucida, MADSEN 113 p. 158 fig. 6.—Peniagone (or S.) sp., MADSEN 113 p. 159 fig. 7.

Stichopus horrens p. 125, variegatus p. 123 with var. hermanni p. 124, Domentay 46.—S. badionotus, macroposentheses, Fontaine 61 p. 31.—S. horrens, Tokioka 218 p. 147 pl. vii figs. 8—11.—S. regalis, Toktorese 220 p. 227 fig. 8.—S. japonicus, tremulus, Turbellaria parasitic in, Westelad 238.

Synapta maculata, Domentay 46 p. 126.

Synaptula hydriformis, Fontaine 61 p. 32 fig.

Thelenota ananas, Domentay 46 p. 126.

Thyone suspecta p. 31, T. sp. p. 31 fig., Fontaine 61.—T. fusus mediterranea, Tortonese 220 p. 233.

ECHINOIDEA

†Radiole undet., Walter 235 p. 701 • pl. lxii fig. 5.

Abatus agassizii p. 46 pl. ix figs. 4–5, pl. xxvi figs. 1–2, 5, pl. xxvi figs. 1–2, 5, pl. xxvii figs. 1–2, 5, pl. xxvii figs. 1–6, pl. xvvii figs. 1–4, cavernosus p. 39 pl. vi figs. 1–7, pl. xviii figs. 1–2, pl. xxiii figs. 1–6, curvidens p. 46, pl. xxii figs. 1–6, pl. xxv fig. 5, pl. xxii figs. 1–6, pl. xxv fig. 5, pl. xxii figs. 1–3, philippii p. 42 pl. xxvi figs. 1–7, shackletoni p. 50, Bernasconi 8.

†Abertella gen. n. for type Scutella aberti Conrad 1842 Miocene U.S.A., DURHAM J. Paleont. 27 pp. 350-351 fig. 1C.

Agassizia scrobiculata, H. L. CLARK 30 p. 339 pl. lxiii figs. 62-63.

†Agassizia caobaensis sp. n. Eccene Cuba, Roig An. Acad. Habana 91 pp. 172-173 pl. xii figs. 1-2.

Alexandria see Clypeaster.

Allocentrotus fragilis, H. L. CLARK 30 p. 276 pl. xliv fig. 21.

Amblypneustes ovum var. pachistus, Darin, Bennett & Pope 40 p. 330 pl. xcii fig. 2.

†Amblypygus depressus sp. n. Eocene Cuba, Rote An. Acad: Habana 91 pp. 144-145 pl. iii fig. 10.—A. sp., Darte-Velle 41 p. 43.

†Amphidetus intermedius pp. 40, 91 pl. viii fig. 6, A. sp. n. indet. pp. 40, 92 pl. ii fig. 8, Szörényi 214.

Amphineustes koehleri p. 35 pl. xix figs. 1-6, lorioli p. 34, similis p. 36, Bernasconi 8.

†Amphiope see Echinodiscus.

Anomalanthus see Clypeaster.

†Anomalanthus gigas sp. n. Oligocene Cuba, Roig Rev. Agric. Habana for 1953 p. 54 pls. i-ii.

†Anorthoscutum Lambert & Thiéry 1914 a synonym of Scutellaster Cragin 1895, Durham 49.

†Antillaster expansus sp. n. Oligocene Cuba pp. 63-64 pl. xviii, jaumei sp. n. Miocene Cuba pp. 64-65 pl. xv pars, Roig Rev. Agric. Habana for 1953.

Apatopygus recens, Fell 55 p. 107.

Arbacia dufresnei, Bernasconi 8 p. 13 pl. v, pl. vi figs. 8-10, pl. vii fig. 4, pl. viii figs. 7-11.—A. incisa p. 244 pl. xl fig. 11, \$patuligera p. 246 pl. xl fig. 12, H. L. Clark 30.—A. lixula africana, Dartevelle 41 p. 26 fig. 5, pl. A fig. 2.—A. lixula, Neefs 151.—A. aequitaberculata, Palombi & Santarelli 155 p. 215 figs.—A. lixula, Tortonese 220 p. 212 fig. 7.

†*Arbacina* ? sp. n., Szörényi **214** pp. 11, 58 pl. vi fig. 7.

†Archaeopneustes of. hystrix, Morishita 141 p. 28 pl. iii figs. 2-3.

†Asterobrissus pomeli, DARTEVELLE 41 p. 45.

Astriclypeus manni, Morishita 139 p. 108 fig. 2b, pl. xi fig. 2.

†Astriclypeus manni minoensis subsp. n. Miocene Japan, Morishitta Mem. Coll. Sci. Kyoto B 20 1952 p. 111 pl. xi fig. 1; A. integer p. 108, A. sp. p. 108 fig. 2a, ID.—A. manni p. 62 pl. vi fig. 3 with subsp. minoensis p. 63 pl. vi figs. 1-2, Morishitta 142.

†Astrodapsis nipponicus, Shikama 182 pl. lv fig. 18.

Astropyga pulvinata, H. L. CLARK 30 p. 236 pl. xxxix fig. 9.

Austrocidaris canaliculata p. 4 pls. ii-iii with var. lorioli p. 6, A. spinulosa p. 6 pl. i, Bernasconi 8.

†Barnumia gen. n. p. 29 with type B. browni sp. n. Cretaceous Guatemala, Cooke Prof. Pap. U.S. geol. Surv. 254A p. 30 pl. xvi figs. 2-5.

Biarritzella see Clypeaster.

Brisaster moseleyi, Bernasconi 8 p. 56 pl. xxxii figs. 1-2.—B. townsendi, H. L. Clark **30** p. 340 pl. lxiv fig. 64.

†Brissopatagus rojasi sp. n. Eocene Cuba, Roig An. Acad. Habana 91 p. 164 pl. ix figs. 5-6.

Brissopsis pacifica, H. L. CLARK 30 p. 341 pl. Ixv figs. 66-67.—B. lyrifera, Iarval stages, Rees 165 figs.—B. lyrifera, Tortonese 220 p. 224.

Brissus latecarinatus, H. L. Clark 30 p. 343 pl. lxvii fig. 74, pl. lxviii fig. 75. —B. brissus, Fontaine 60 p. 8.—B. unicolor, Tortonese 220 p. 224.

†Brissus caobaense sp. n. Eocene Cuba, Roig An. Acad. Habana 91 p. 163 pl. vii fig. 5, pl. viii figs. 1-2.— B. jacquementi, Szörényi 214 pp. 41, 92 pl. iv fig. 5.

Bunactis see Clypeaster.

Caenocentrotus gibbosus, H. L. CLARK 30 p. 265 pl. xliii fig. 20.

†Cardiaster deciper sp. n. Cretaceous U.S.A., COOKE Prof. Pap. U.S. geol. Surv. 254A pp. 27–28 pl. x figs. 3–6; cuttus p. 28 pl. xi figs. 1-7, marylandicus p. 28 pl. xi figs. 1-7, marylandicus p. 28 pl. xiv figs. 1-5, smocki p. 28 pl. xi figs. 8–10, In.—C. kelleri, Dartevelle 41 p. 135 figs. 41–42.

Cassidulus pacificus, H. L. CLARK 30 p. 338 pl. lxii fig. 61.

†Cassidulus rojasi sp. n. Oligocene Cuba, Roig An. Acad. Habana 91 pp. 146-147 pl. iv figs. 3-5.

†Catopygus mississippiensis sp. n. Cretaceous U.S.A., Cooke Prof. Pap. U.S. geol. Surv. 254A pp. 15-16 pl. v figs. 3-6; pusillus, ID. p. 15 pl. v figs. 1-2.

†Catopygus (Oolopygus) gonzalezi sp. n. Cretaceous Cuba, Roig An. Acad. Habana 91 pp. 145-146 pl. iv figs. 1-2. —C. (O.) williamsoni, Cooke 33 p. 16 pl. v figs. 7-10.

Centrechinus mexicanus, H. L. Clark 30 p. 235 pl. xxxviii fig. 8.—C. antillarum, Fontaine 60 p. 4 fig.

Centrocidasis doederleini, H. L. CLARK 30 p. 234 pl. xxxviii fig. 7.

Centrostephanus coronatus, H. L. Clark 30 pp. 237, 244 pl. xxxix fig. 10.—C. rodgersii, Dakin, Bennett & Pope 40 p. 329 pl. xci fig. 3.—C. longispinus, Tortonese 220 p. 214.

†Cheopsia mortenseni, Dartevelle 41 p. 154 pl. xii figs. 10, 13-14.

Cidaris cidaris, Tortonese 220 p. 210.

†Cidaris pyrenaica, Bataller 6 p. 79.—C. texanus p. 5 pl. i fig. 14, wahalakensis p. 5 pl. i figs. 1-2, Cooke 33.—C. malheiroi p. 5, vafellus p. 6 pl. A fig. 1, Dartevelle 41.—C. coronota, fig. 8a, C. sp. [Radiole] fig. 8b, Kuhn-Schnyder 96 p. 216.—C. galeotti p. 20 pl. i figs. 1-7, mullerriedi p. 22 pl. i figs. 8-9, submarginata p. 23, texanus p. 24 pl. i figs. 10-11, sp. A p. 25 pl. i figs. 12-16, sp. B p. 25 pl. i figs. 17-18, Maldonado-Koerdell 115.—C. herthae, Müller 147 p. 26 pl. i fig. 1.

†Clarkiella hemispherica, Cooke 33 p. 18 pl. v figs. 15-18.

Clypeaster elongatus sp. n. Galapagos Is., H. L. Clark Allan Hancock Pacif. Exped. 8 1948 pp. 308–309 pl. xlviii fig. 30, pl. xlix fig. 31, pl. l fig. 33; europacificus p. 294 pl. xlvi fig. 26, ochrus p. 297 pl. xlvii fig. 27, rotundus p. 296 pl. xlvii fig. 28, speciosus p. 308 pl. xlviii fig. 29, Ib.—C. rosaceus p. 6 fig., subdepressus p. 7 fig., Fontaine 60.

—C. with subgenera and sections Alexandria, Anomalanthus, Biarritzella, Bunactis, Clypeaster, Coronanthus, Guebhardanthus, Laganidea, Laubeanthus, Leptoclypus, Orthanthus, Oxyclypeina, Paleanthus, Paratinanthus, Pavaya, Platyclypeina, Pliophyma, Rhaphidoclypus, Stolonoclypus, Tholeopelta, validity discussed, and recent and fossil spp. assigned, ROMAN 173.

†Clypeaster lopezriosi sp. n. Oligocene Cuba, Rotg An. Acad. Habana 91 p. 139 pl. i fig. 6.—C. julii nom. n. for Anomalanthus gregoryi Lambert 1915 Miocene West Indies, non Clypeaster gregoryi Lambert 1913, Roman Bull. Soc. géol. Er. 6 2 p. 408 footnote 2.—C., discussion of African fossil spp., Dartevelle 41 p. 69; C. sp., Io. p. 60 fig. 12, pl. xv figs. 4–5.—C. parischi, Szörényi 214 pp. 20, 68 pl. ii fig. 3.

Clypeaster (Bunactis) rangianus, DARTEVELLE 41 p. 66 pl. xiii fig. 1.

†Clypeaster (Bunactis) rangianus p. 68 pl. B figs. 1-2, C. (B.) sp. p. 61 fig. 13, pl. viii figs. 1-2, Dartevelle 41.

Clypeaster (Clypeaster) rosaceus, Roman 173 p. 404 fig. C.

Clypeaster (Rhaphidoclypus) reticulatus, Dartevelle 41 p. 68 pl. xiii fig. 2.—C. (R.) reticulatus, Roman 173 p. 404 fig. C.

†Clypeaster (Rhaphidoclypus) armadilloensis sp. n. Oligoeene Cuba, Rote An. Acad. Habana 91 pp. 139–140 pl. ii figs. 1–2.—C. borgesi, Darteville 41 p. 64 pl. xiv figs. 1–2.

Clypeaster (Stolonoclypus) humilis p. 404 fig. Ba, †virescens p. 404 fig. Bb, Roman 173.

†Clypeus plotii, Sikošek 187 p. 152 pl. i fig. 1.

†Codiopsis stephensoni sp. n. Cretaceous Texas, Cooke Prof. Paper U.S. geol. Surv. 254A p. 8 pl. i figs. 9-11.—C. meslei, Groove 63 p. 351.

Colobocentrotus mertensii, Ohshima 153 p. 160 fig. 2, pl. i fig. 4, pl. ii figs. 10-11.

†Conoclypeus pinfoldi sp. n. Eocene Pakistan, GLL J. Paleont. 27 pp. 843– 844 pl. 91 figs. 1–3; pilgrimi, Id. p. 843 pl. 91 fig. 4.

†Conulus stephensoni sp. n. Cretaceous Texas, Cooke Prof. Pap. U.S. geol. Surv. 254A p. 11 pl. iii figs. 8-13.

Coronanthus see Clypeaster.

†Cottaldia aff. benettiae, Dartevelle 41 p. 28.

†Crucibrissus cabrerai sp. n. Eocene Cuba, Roig Rev. Agric. Habana for 1953 p. 67 pl. x pars, pl. xxi.

Ctenocidaris geliberti p. 8, perrieri p. 9, speciosa p. 8 pl. iv, Bernasconi 8.

†Cylindrolampas see Echinolampas,

 $\dagger Cypholampas$ see Echinolampas.

†Deakia armadilloensis sp. n. Eocene Cuba, Roig An. Acad. Habana 91 pp. 165-166 pl. x figs. 1-2.

Dendraster laevis sp. n. California & N.W. Mexico pp. 312-313 pl. 1 figs. 34-36, mexicanus sp. n. Lower California pp. 313, 317 pl. li figs. 37-39, rugosus sp. n. Lower California pp. 318-319 pl. lii figs. 40-41, H. L. CLARK Allan Hancock Pacif. Exped. 8 1948; excentricus p. 310 pl. xix fig. 32, In.

†Dendraster ashleyi p. 68 pl. xvii figs. 11-12, pl. xviii figs. 12-14, pl. xix fig. 6, pl. xx figs. 1, 5-6, cf. coalingaensis p. 67 pl. vii figs. 6, 8, Woodring & Bramlette 241.

Diadema antillarum, MILLOTT 123 p. 497 figs. 1-11, 125 p. 98 figs. 1-2, 127 p. 973 figs. 1-3 [iridophores].—D. antillarum, Turbellaria parasitic in, Westblad 238.

†Ditremaster olbrechtsi sp. n. Eocene Congo & Angola, Dartevelle Ann. Mus. Congo belge 13 pp. 148-153 figs. 48-50, pl. xi figs. 1-5.

† Douvillaster benguellensis, DARTE-VELLE 41 p. 144 figs. 45-46.

†Echinarachnius rumoensis sp. n. Miocene Japan, Науазака & Shibata J. Fac. Sci. Hokkaido Univ. Geol. Min. 8 1952 pp. 82-85 figs. 1 а-с.—Е. падапоепзіз sp. n. Pliocene Japan, Morishita Mem. Coll. Sci. Kyoto В 20 pp. 220-222 fig. 3a, pl. i fig. 4; microthyroides p. 219 pl. i fig. 3, parma obesus p. 221 fig. 3d, E. rumoensis p. 221 fig. 3c, subtumidus p. 221 fig. 3b, ID.—E. mirabilis, Shikama 182 pl. xliv fig. 14.

†Echinarachnius (Kewia) nipponica, Shikama 182 pl. xxxii fig. 17.

Echinocardium cordatum, Dakin, Bennett & Pope 40 p. 332 pl. sciii fig. 1.—E. cordatum, Lafon 100 p. 152.
—E. cordatum, flavescens, larval stages, Rees 165 figs.—E. mortenseni, Tortonese 219 p. 271.—E. cordatum p. 221,

flavescens p. 222, mediterraneum p. 222, mortenseni p. 223 pl. i fig. 9, TORTONESE 220.—E. flavescens, Turbellaria parasitic in, Westelad 238 p. 271.

†Echinoconus castanea, Maksimović & Marković 114 p. 186 pl. i figs. 4-5.

†Echinocorys texanus, Cooke **33** p. 25 pl. x figs, 1-2, pl. xi fig. 17.—E. sp., MÜLLER **147** p. 32 pl. i fig. 3.—E. scutata, WILCOX **239** p. 83 figs. 1-6.

†Echinocrinus sp., Chronic 27 p. 165 pl. xxxv figs. 13-17.

Echinocyamus pusillus, Tortonese 220 p. 220.

†Echinocyamus crispus, Morishita 142 p. 61 pl. vi figs. 4-5.

Echinodiscus bisperforatus, DARTE-WELLE 41 p. 74 fig. 14, pl. xv figs. 1-2.

†Echinodiscus chikuzenensis, Mori-Shita 139 p. 107 fig. 2c.

†Echinodiscus (Amphiope) neuparthi, Dartevelle 41 p. 75 figs. 15-16, pl. viii tigs. 4, 6, pl. xv fig. 3.

†Echinolampas consolationis sp. n. Miocene Cuba, Roig An. Acad. Habana '91 pp. 148-149 pl. iv figs. 9-11; brachyona, Id. p. 148.—E. atrophus podolicus
subsp. n. Miocene Ukraine, Szörényi
Geol. hung. Ser. Palaeont. 23 pp. 34,
35 pl. vi fig. 4; E. barcinensis pp. 34,
35, hemisphaericus pp. 34, 84 pl. iii
fig. 2, Id.—E. angustipetalus p. 143 pl.
xii fig. 2, dumasi p. 143 pl. xiii fig. 3,
'Dechović & Seneš 19.—E. spp.,
Dartevelle 41 p. 57 pl. vi figs. 1-3
en bas.—E. angulatus p. 174 pl. i fig. 1,
vyramidalis p. 174 pl. i fig. 2, Pinar
& Roman 163.—E. yoshiwarai, Shikama
182 pl. xli fig. 4.

†Echinolampas (Cylindrolampas) sandiegensis sp. n. Eccene Cuba, Rote An. Acad. Habana 91 pp. 151-152 pl. v igs. 3-4; altiesima, Id. p. 150.

†Echinolampas (Cypholampas) eocenirus sp. n. Eocene Cuba, Roig An. Acad. Habana 91 p. 149 pl. v figs. 1-2.

†Echinolampas (Heteroclypeus) hungvricus, Čechovič & Seneš 19 p. 143 bl. xii fig. 1, pl. xiii fig. 1.

†Echinolampas (Macrolampas) gigas 19. n. Miocene Cuba, Rote Rev. Agric. Habana for 1953 pp. 55-56 pls. iii pars), vii-viii. †Echinolampas (Palaeolampas) acuminatus p. 175 pl. i fig. 3, aff. francei p. 178 fig. 1, pl. i fig. 4, Pinar & Roman 163.

†Echinolampas (Progonolampas) moronensis pp. 152–153 pl. vi figs. 1–2, torrense p. 152 pl. v figs. 5–6, spp. n. Oligocene Cuba, Roig An. Acad. Habana 91.

Echinometra oblonga p. 281 pl. xlv fig. 24, vanbrunti p. 293 pl. xlvi fig. 25, H. L. CLARK 30.—E. lacunter, DARTEVELLE 41 p. 38 fig. 6, pl. A fig. 6, pl. i figs. 7-8.—E. lucunter p. 5 figs., viridis p. 6 fig., FONTAINE 60.

†Echinometra lacunter, Dartevelle 41 p. 38 figs. 7-8, pl. A fig. 5, pl. i figs. 4-6.

Echinoneus cyclostomus, Fontaine 60 p. 7 figs.

†Echinoneus robustus sp. n. Oligocene Cuba, Roig An. Acad. Habana 91 p. 140 pl. iii figs. 1-2.

†Echinopsis friryi, Dartevelle 41 p. 33 pl. iii figs. 3-15, 17-20.

†Echinospatangus sp. indet., HASSAN 68 p. 128.

Echinus acutus p. 215 pl. i fig. 8, melo p. 217, Tortonese 220.

Encope emarginata, Bernasconi 8 p. 27 pl. xiii, pl. xvii figs. 8-11, pl. xxxii fig. 3.—Encope p. 319 with E. cocosi sp. n. Cocos Id. p. 330 pl. lvi figs. 48-49, ecuadorensis sp. n. Ecuador pp. 333-334 pl. lix figs. 54-55, fragilis sp. n. Mexico pp. 335-336 pl. lx figs. 56-57, insularis sp. n. Mexico pp. 336-337 pl. lxi figs. 56-59, irregularis sp. n. Colombia to Costa Rica pp. 332-333 pl. lviii figs. 52-53, laevis sp. n. Nicaragua pp. 327-328 pl. liv fig. 45, pl. lv fig. 46, H. L. CLARK Allan Hancock Pacif. Exped. 8 1948; E. galapagensis p. 331 pl. lvii figs. 50-51, grandis p. 325 pl. liii fig. 42, micropora p. 329 pl. lv fig. 47, perspectiva p. 326 pl. liii fig. 43, wetmorei p. 326 pl. liv fig. 44, ID., text-figs. 1-2.

†Eoscutum see Fibularia.

†Epiaster carvalhoi sp. n. Cretaceous Angola, Dartsvelle Ann. Mus. Congo belge 13 pp. 142-143 pl. x figs. 1-2; angolensis p. 141, catumbellensis p. 137 figs. 43-44, pl. x figs. 3-4, Id.

Eucidaris thouarsii, H. L. Clark 30 p. 229 pl. xxxv fig. 1.—E. tribuloides africana, Dartevelle 41 p. 8 fig. 1, pl. A fig. 3.—E. tribuloides, Fontaine 60 p. 4 fig.

†Eucidaris tribuloides africana, Dartevelle 41 p. 8 pl. A fig. 4, pl. i figs. I-3.

†Eupatagus brodermanni pp. 159-160 pl. ix figs. 1, 4, calistoides pp. 160-161 pl. vi fig. 8, casanovai p. 159 pl. ix figs. 2-3, turibacoensis pp. 157-158 pl. viii figs. 3-4, spp. n. Eocene Cuba, Roig An. Acad. Habana 91; alatus p. 156, attenuatus p. 156, ocalanus p. 161, ID.

†Eupatagus (Perispatangus) daradensis, Dartevelle 41 p. 165 fig. 53, pl. xi figs. 6-8.

†Eupatagus (Spatangomorpha) pinarensis sp. n. Eocene Cuba, Rore An. Acad. Habana 91 p. 157 pl. vii figs. 3-4.

Faorina chinensis, HAYASAKA 71 p. 217 pl. xiv figs. 1-6.

†Faujasia chelonium sp. n. Cretaceous Texas, Cooke Prof. Pap. U.S. geol. Surv. 254A p. 14 pl. iv figs. 11-14.

†Fibularia sandalina sp. n. Miocene Ukraine, Szörényi Geol. hung. Ser. Palaeont. 23 pp. 13, 60 pl. v fig. 6.

 $\dagger Fibularia$ (Eoscutum) seefriedi, Cox 36 p. 35 pl. v figs. 17–20.

Fibularia (Fibulariella) acuta, To-KIOKA 218 p. 146 figs. 1-2.

Genocidaris maculata, Tortonese 220 p. 214.

†Globator vaughani sp. n. Cretaceous Texas, Cooke Prof. Pap. U.S. geol. Surv. 254A p. 12 pl. iv figs. 1-4.

Goniocidaris maculata, Tortonese 219 p. 271.

Goniomaretia laevis, H. L. Clark 30 p. 347 pl. lxx figs. 80-81.

†Goniopygus zitteli telostocensis subsp. n. Cretaceous Mexico, Maldonado-Koerdell An. Escu. nac. Cienc. biol. Méx. 7 pp. 38–39 pl. ii figs. 38–40; G. bolaensis, Id. p. 38.

Guebhardanthus see Clypeaster.

†Haimea cylindrica p. 141 pl. iii figs. 6–7, gigantea p. 143 pl. iii fig. 11, pentagona pp. 142–143 pl. iii figs. 8–9, subcylindrica pp. 141–142 pl. iii figs. 4-5, spp. n. Eocene Cuba, Rote An. Acad. Habana 91; caillaudi, In. p. 143 pl. iii fig. 3.

†Hardouinia mcglameryae sp. n. Cretaceous U.S.A., COOKE Prof. Pap. U.S. geol. Surv. 254A p. 24 pl. viii figs. 23-26; aequorea p. 22 pl. viii figs. 15-22, bassleri p. 22 pl. viii figs. 6-11, florealis p. 23 pl. viii figs. 11-14, kellumi p. 21 pl. vii figs. 1-5, micrococcus p. 21 pl. vi figs. 1-10, pl. vii fig. 12, mortonis p. 19 pl. v figs. 19-21 with var. emmonsi p. 20 pl. v figs. 22-24, H. potosiensis p. 19, stantoni p. 23 pl. viii figs. 7-10, taylori p. 24 pl. viii figs. 1-6, In.

Heliocidaris erythrogramma p. 331 pl. lxxxix, tuberculata p. 332 pl. xc, Dakin, Bennett & Pope 40.

†Hemiaster amelianus sp. n. Cretaceous Guatemala p. 35 pl. xvi fig. 1, arcolensis sp. n. Cretaceous Alabama pp. 32-33 pl. xii figs. 12-16, sabinal sp. n. Cretaceous Texas p. 34 pl. xiii figs. 7-10, Cooke Prof. Pap. U.S. geol. Surv. 254A; beecheri p. 31, delawarensis p. 35 pl. xiv figs. 6-9, humphreysanus p. 30 pl. xiii figs. 1-4, jacksoni p. 33 pl. xiii figs. 5-11, "H." kümmeli p. 35 pl. xiii figs. 5-11, "H." kümmeli p. 35 pl. xiii figs. 5-1, "H. iii figs. 10-14, unqula p. 34 pl. xiv figs. 10-14, welleri p. 35 pl. xiv figs. 15-17, wetherbyi p. 31 pl. xii figs. 17-18, ID.—H. gonzalezmuñozi sp. n. Cretaceous Cuba, Roig An. Acad. Habana 91 pp. 170-171 pl. xi figs. 3-4.—H. reineckei p. 146, H.? sp. p. 146 fig. 47, pl. iv figs. 5-6, Dartevelle 41.—H. chauveneti, meslei, Gigout 63 p. 362.

†Hemicidaris villadai sp. n. Cretaceous Mexico, MALDONADO-KOERDELL An. Escu. nac. Cienc. biol. Méx. 7 pp. 27-28 pl. i figs. 22-24.

†Hemicidaris (Pseudocidaris) clunifera, Maldonado-Koerdell 115 p. 26 pl. i figs. 19-21.

Hemiheliopsis fonti, Dartevelle 41 p. 85 figs. 17-18.

†Hemipatagus sp., HASSAN 69 p. 15 pl. A fig. 1.

Hesperocidaris asteriscus sp. n. Panama, H. L. Clark Allan Hancock Pacif. Exped. 8 1948 pp. 233-234 pl. xxxvi fig. 4, pl. xxxvi figs. 5-6; panamensis p. 230 pl. xxxv fig. 2, perplexa p. 231 pl. xxxvi fig. 3, ID.

†Hessotiara zuberti sp. n. Jurassic Switzerland, Jeannet Vjschr. naturf. Ges. Zürich 98 pp. 176-177 figs. 1-5.

†Heteroclypeus see Echinolampas.

†Heterosalenia alloiteaui sp. n. Jurassie N. Africa, ZOEKE Bull. Soc. géol. Fr. 6 2 pp. 249-252 figs. 1-2, pl. xib figs. 1-3.

†Holaster feralis sp. n. Cretaceous U.S.A., Cooke Prof. Pap. U.S. geol. Surv. 254A pp. 26-27 pl. ix figs. 1-5, pl. x fig. 7.—H. lerichei sp. n. Cretaceous Angola, Dartevelle Ann. Mus. Congo belge 13 pp. 131-135 fig. 40, pl. ix figs. 1-4; dombeensis, Id. p. 130.—H. laevis, Marsimović & Marković 114 p. 186.

†Holectypus montalvensis sp. n. Cretaceous Cuba, Roig An. Acad. Habana 91 p. 138 pl. i figs. 2-3.—H. sp., Darrevelle 41 p. 186 fig. 57, pl. xix figs. 1-12.

Holopneustes pycnotilus, Dakin, Bennett & Pope **40** p. 330 pl. xeii fig. 1.

†Hypsoclypus fuentesi p. 55 pl. v, pl. vi (pars), holguinensis pp. 54-55 pls. iii-iv (pars) spp. n. Miocene Cuba, Roig Rev. Agric. Habana for 1953.—H. doma p. 179, montesiensis p. 178 pl. i fig. 5, Pinar & Roman 163.

†Hypsoheteroclypus nom. n. for type Hypsoclypus doma Pomel 1887 Miocene Algiers, Szörényi Geol. hung. Ser. Palaeont. 23; vicinoconoideus sp. n. pp. 33, 83 pl. vi figs. 2-3, plagiosomus corsicarus subsp. n. pp. 28, 78 pl. iv fig. 1, Miocene Europe, In.; H. acuminatus pp. 32, 82 pl. vi fig. 1, hungaricus pp. 31, 81 pl. iii fig. 3, montesiensis pp. 30, 80 pl. v fig. 3, plagiosomus with subsp. lamberti pp. 30, 80 pl. v figs. 1-2 & subpentagonalis pp. 29, 78 pl. iv fig. 2, H. pyramidalis pp. 32, 83 pl. ii fig. 4, In.

Idiobryssus coelus, H. L. Clark 30 p. 343 pl. lxvii figs. 71-73.

†Isomicraster danei p. 29 pl. xi figs. 11-13, rossi p. 29 pl. xi figs. 15-16, spp. n. Cretaceous U.S.A., COOKE Prof. Pap. U.S. geol. Surv. 254A.

Kewia see Echinarachnius.

Laganidea see Clypeaster.

†Lajanaster rojasi sp. n. Oligocene †Cuba, Roio Rev. Agric. Habana for 1953 p. 58 pl. x pars. †Lambertona gen. n. for type Victoriaster lamberti Roig Eccene Cuba, Roig Mem. Soc. cubana Hist. nat. 21 p. 267 pls. xxvii-xxviii.

†Lanieria uvaldana sp. n. Cretaceous Texas, Cooke Prof. Pap. U.S. geol. Surv. 254A p. 10 pl. iii figs. 1-3.

Laubeanthus see Clypeaster.

†Lefortia trojana sp. n. Cretaceous U.S.A., COOKE Prof. Pap. U.S. geol. Surv. 254A p. 17 pl. iv figs. 8-10.

 $\dagger Leiocidaris$ capelloi, Dartevelle 41 p. 13.

Leptoclypus see Clypeaster.

†Linthia yessoensis sp. n. Palaeogene Japan, Minato J. geol. Soc. Japan 56 1950 pp. 158-159 figs. 1-2.—L. variabilis, Cooke 33 p. 36 pl. xiv figs. 18-25.—L. nipponica, Morishita 140 p. 222 pl. i figs. 5-7.—L. nipponica pl. xili fig. 3, praenipponica pl. xxxii fig. 16, Shirama 182.

†Loriola picteti, Maldonado-Koerdell 115 p. 29 pl. ii figs. 28-30.

Lovenia cordiformis, H. L. CLARK 30 p. 347 pl. lxxi figs. 82-83.

Loxechinus albus, Bernasconi 8 p. 23 pl. vii fig. 1, pl. xi, pl. xii figs. 1-9.— L. albus, H. L. Clark 30 p. 265 pl. xliii fig. 19.

Lytechinus anamesus p. 247 pl. xli fig. 14, pictus p. 249 pl. xli fig. 15, semituberculatus p. 261 pl. xli fig. 16, H. L. Clark 30.—L. variegatus, Fontaine 60 p. 5 fig.—L. semituberculatus, Tortonese 221 p. 479.

 $\dagger Macrolampas$ see Echinolampas.

†Macropneustes brodermanni sp. n. Oligocene? Cuba pp. 62-63 pl. xv pars, pl. xvi pars, gómez-mazue sp. n. Oligocene Cuba p. 63 pl. xvii, Roig Rev. Agric. Habana for 1953.—M. cf. baylei?, Dartevelle 41 p. 136.

†Mauritanaster marroquinensis sp. n. Oligocene Cuba, Roig Rev. Agric. Habana for 1953 pp. 61-62 pl. xiv.

†Meyapatagus gen. n. p. 58 for type M. franciscanus sp. n. Oligocene Cuba, Roic Rev. Agric. Habana for 1953 pp. 59-60 pl. vi pars, pl. xi, depressus sp. n. Oligocene Cuba p. 60 pl. xii, turibacoensis sp. n. Eocene Cuba p. 61 pl. xiii, In.

Mellita platensis p. 31 pl. xvi figs. 1-7, pl. xvii figs. 1-7, quinquiesperforata latiambulacra p. 29 pl. ix fig. 7, pl. xiv, pl. xv figs. 1-7, pl. xxviii figs. 3-4, BERNASCONI 8.—M. longifissa, H. L. Clark 30 p. 337 pl. lxii fig. 60.—M. quinquesperforata fig., sexiesperforata, FONTAINE 60 p. 7.

Meoma grandis, H. L. Clark 30 p. 344 pl. lxviii figs. 76-77.—M. ventricosa,

FONTAINE 60 p. 8 figs.

†Merriamaster cf. perrini, Woodring & Bramlette 241 p. 68 pl. x figs. 11-14.

Mespilia globulus, Оняніма 153 р. 159 pl. ii figs. 5-9.

†Metalia palmeri sp. n. Oligocene Cuba, Rong An. Acad. Habana 91 pp. 166-167 pl. x fig. 3; M. sp., In. p. 167 pl. x fig. 4.

†Micraster uddeni sp. n. Cretaceous Texas, Cooke Prof. Pap. U.S. geol. Surv. 254A pp. 37-38 pl. xv figs. 6-9; americanus, Id. p. 38 pl. xv figs. 10-13. —M.? sp. indet., Hassan 68 p. 129.

Moira clotho, H. L. CLARK 30 p. 341 pl. lxiv fig. 65.—M. atropos, Fontaine 60 p. 8.

†Moira obesa, Shikama 182 pl. xli fig. 5.

†Neocatopygus caobaense sp. n. Eocene Cuba, Roig An. Acad. Habana 91 p. 154 pl. vi figs. 3-4.

†Neopatagus gen. n. for type Breynia cubensis Cotteau 1875 Oligocene Cuba, Roig Mem. Soc. cubana Hist. nat. 21 p. 258; lajanensis p. 260 pl. xxix (upper fig.), stefaninii p. 261 pl. xxix (lower fig.), ID.

Notocidaris mortenseni, Bernasconi 8 p. 10.

†Nucleopygus atlanticus brevior subsp. n. Cretaceous Belgian Congo, Dartevelle Ann. Mus. Congo belge 13 pp. 51-52 pl. v figs. 2, 4-5, 10-11, 13; N. atlanticus, Id. p. 46 figs. 9-11, pl. iv figs. 1-4, 7-11, pl. v figs. 1, 3, 6-9, 12, 14-16.—N. texanus, Cooke 33 p. 13 pl. iv figs. 5-7.

†Oligopigus herrerai pp. 154-155 pl. vi figs. 6-7, 9, sanjosephi pp. 155-156 pl. vii figs. 1-2, spp. n. Eocene Cuba, Roig An. Acad. Habana 91.

†Oolopygus see Catopygus.

†Opissaster cantabriae sp. n. Cretaceous Cuba, Roig An. Acad. Habana 91 pp. 171-172 pl. xi figs. 5-6.

Orthanthus see Clypeaster.

†Orthopsis aguilerai sp. n. Cretaceous Mexico, Maldonado-Koerdall An. Escu. nac. Cienc. biol. Méx. 7 pp. 28–29 pl. i figs. 25–27.—O. cf. ruppellii p. 28, O. sp. p. 30, Dartevelle 41.

Ova see Schizaster.

Oxyclypeina see Clypeaster.

†Palaeechinus canadensis sp. n. Carboniferous Canada, Kier Geol. Mag., London 90 pp. 65-69 figs. 1-4.

†Palaeolampas see Echinolampas.

†Palaeolampas elongatus sp. n. Eocene Cuba, Rom Rev. Agric. Habana for 1953 pp. 56–57 pl. iv (pars).—P. alta, plateia, Rom 170 p. 150.

†Palaeopneustes cf. cristatus, Morishita 141 p. 27 pl. iii fig. l.

Paleanthus see Clypeaster.

Paracentrotus lividus, Palombi & Santarelli 155 p. 214 figs.—P. lividus, Tortonese 220 p. 218.—P. lividus, Ubisch 227 p. 294 fig. I [Pluteus].

†Paralampas conceptionis sp. n. Eccene Cuba, Roig An. Acad. Habana 91 pp. 153-154 pl. vi fig. 5.

Parapneustes cordatus p. 36, reductus p. 37 pl. ix fig. 6, pl. xx figs. 1-3, pl. xxi figs. 7-10, Bernasconi 8.

Paratinanthus see Clypeaster.

†Parascutella gen. n. for type Scutella leogranessis Lambert 1903 Miocene France, Durham J. Paleont. 27 pp. 349-350 fig. 1A.

†Paraster delgadoi sp. n. Cuba p. 174 pl. xii figs. 5-6, pinarensis sp. n. Ecocene Cuba p. 173 pl. xii figs. 3-4, Roig An. Acad. Habana 91.

† Parmulechinus, diagnosis, Durham 50 pp. 349, 351.

Pavaya see Clypeaster.

†Pedinopsis pondi, COOKE 33 p. 7 pl. ii figs. 1-5.

† Periaster sp., Dartevelle 41 p. 157.

†Pericosmus artemisae sp. n. Miocene Cuba pp. 168–169 pl. xi fig. l, valenzuelai sp. n. Miocene Cuba pp. 167–168 pl. x fig. 5, zanolettii sp. n. Eocene Cuba pp. 169–170 pl. xi fig. 2, Rotg An. Acad. Habana 91.—P. delgadoi sp. n. Oligocene Cuba, Rotg Rev. Agric. Habana for 1953 p. 65 pl. xvi pars. †Perispatangus see Eupatagus.

Phyllacanthus parvispinus, Dakin, Bennett & Pope 40 p. 329 pl. xci figs. 1-2.

†Phyllobrissus cubensis, Cooke 33 p. 17 pl. v figs. 11-14.

†Phymosoma gigantea sp. n. Eocene Cuba, Roig An. Acad. Habana 91 p. 136 pl. ii fig. 1.—P. hilli, Cooke 33 p. 9 pl. iii figs. 6-7.—P. sp., Dartevelle 41 p. 18 figs. 3-4, pl. ii figs. 1-8.—P. aquitanicum p. 33, mexicanum p. 34, aff. mexicanum p. 35, P. texanum p. 36 pl. ii figs. 36-37, volanum p. 37, Maldonado-Koerdell 115.

Plagiobrissus pacificus, H. L. CLARK 30 p. 342 pl. lxvi figs. 68-70.—P. africanus, Dartevelle 41 p. 167 fig. 54.

†Plagiobrissus sp., Dartevelle 41 p. 169 fig. 56, pl. viii fig. 5.—P. loveni, robustus, Roig 170 p. 162.

†Planilampas circularis sp. n. Oligocene Cuba, Rom Rev. Agric. Habana for 1953 p. 57 pl. ix.

Platyclypeina see Clypeaster.

Plexechinus nordenskjöldi, Bernasconi 8 p. 57.

†Pliolampas sp., Dartevelle 41 p. 54 pl. vi figs. 1-3 en haut.—P. gauthieri pp. 21, 70 pl. v fig. 5, vassalli pp. 21, 69 pl. iii fig. 5, Szörényi 214.

Pliophyma see Clypeaster.

†Porocidaris lopezi sp. n. Eocene Cuba, Roig An. Acad. Habana 91 p. 137 pl. i fig. 4.

†Porosoma reesidei sp. n. Cretaceous U.S.A., COOKE Prof. Paper U.S. geol. Surv. 254A pp. 6-7 pl. i figs. 5-8.

Pourtalesia miranda, Mortensen 143 p. 47 fig. 1.

†Procassidulus echevarriai sp. n. Oligocene Cuba, Roig An. Acad. Habana 91 p. 145 pl. iv fig. 8.

Progonolampas see Echinolampas.

†Proraster dalli, Clarke 33 p. 36 pl. xv figs. 1-5.

†Prospatangus acuminatus sp. n. Miocene Ukraine, Szönényi Geol. hung. Ser. Palaeont. 23 pp. 39, 91 pl. vii fig. 5; austriacus pp. 37, 88 pl. viii fig. 3, corsicus pp. 36, 87 pl. vii fig. 3, delphinus pp. 35, 86 pl. vii figs. 1–2, fabianii pp. 39, 90 pl. viii fig. 5, hungaricus pp. 38, 89

pl. viii fig. 4, pustulosus pp. 37, 88 pl. vii fig. 4, P. sp. n. pp. 39, 91 pl. vii fig. 6,

Psammechinus microtuberculatus, Palombi & Santarelli 155 p. 215 figs.— P. microtuberculatus, Tortonese 220 p. 217.

†Psammechinus cf. dubius, DARTE-VELLE 41 p. 36 pl. i figs. 9-11.

†Pseudananchys stephensoni sp. n. Cretaceous Texas, Cooke Prof. Pap. U.S. Geol. Surv. 254A p. 26 pl. ix figs. 6-9.

Pseudechinus magellanicus, BERNA-SONI 8 p. 17 pl. vii figs. 2-3, pl. viii figs. 1-6, pl. xviii figs. 3-4.—P. novaezelandiae, FELL 55 p. 105.

†Pseudoastrodapsis nom. n. for Nipponaster Durham 1952 non Niponaster Lambert 1920, Durham J. Paleont. 27 p. 756.

Pseudocidaris see Hemicidaris.

†Pseudorthopsis rojasi sp. n. Eocene Cuba, Roig An. Acad. Habana 91 pp. 137-138 pl. i fig. 5.

†Pygurostoma geometricum p. 14 pl. iv figs. 15-20, pasionense p. 14 pl. xvi figs. 6-11, Cooke 33.

† Pygurus africanus, Dartevelle 41 p. 42.

†Rachiosoma hondoensis sp. n. Cretaceous Texas, Cooke Prof. Paper U.S. geol. Surv. 254A pp. 8-9 pl. iii figs. 4-5; mortoni, Ib. p. 9 pl. i figs. 12-13.

Radiorotula orbiculus with subspp. dentata, radiata semisol, solaris, Darte-Velle 41 p. 108 figs. 24-28, 30-31, pl. C, pl. xvii figs. 1-9.

†Radiorotula orbiculus p. 108 figs. 17-18, 29 with subsp. radiata semisol p. 108 pl. xvii fig. 10, Dartevelle 41.

Rhaphidoclypus see Clypeaster.

†Rhenechinus hopstätteri gen. n. sp. n. Devonian Germany, Dehm Notizbl. hess. Landesamt. Bodenforsch. 81 pp. 88-95 figs. 1-2, pl. v.

Rotula angusti, Dartevelle 41 p. 118 figs. 32-36, pl. xviii figs. 1-4.

†Rotula cf. angusti, Dartevelle 41 p. 123 figs. 37-39, pl. viii fig. 3, pl. xviii figs. 5-7.

Rotulinae, discussion of distribution of fossil and recent spp., DARTEVELLE 41 p. 129.

†Rotuloidea vieirai sp. n. Miocene Luanda, Dartevelle Ann. Mus. Congo belge 13 pp. 103-107 figs. 22-23, pl. vi fig. 4, pl. vii figs. 1-2, pl. xvi figs. 1-8; fimbriata, Id. p. 100 figs. 19-21, pl. xvi figs. 9-14.

†Salenia hondoensis sp. n. Cretaceous Texas, Cooke Prof. Paper U.S. geol. Surv. 254A p. 6 pl. i figs. 3-4.—S. dombeensis p. 15 with subsp. triangularis p. 16 fig. 2, Dartevelle 41.—S. prestensis mexicana p. 39 pl. ii figs. 41-43, S. sp. p. 41, Maldonado-Koerdell 115.

†Schizaster kinasaensis sp. n. Pliocene Japan, Morishita Mem. Coll. Sci. Kyoto B20 pp. 224–225 pl. i fig. 8; S. sp., Id. p. 226.—S. gigas p. 66 pl. xx, pentagonalis pp. 65–66 pl. xix, spp. n. Eocene Cuda, Roig Rev. Agric. Habana for 1953.—S. karreri, Čechovič & Seneš 19 p. 139 pl. xiii fig. 2.—S. douvillei, Dartevelle 41 p. 159 figs. 51–52 pl. xii figs. 1–9, 11–12, 15–17.—S. sp. ind. p. 294 pl. i fig. 10, Köster 93.—S. sp. indet., Morishita 142 p. 63 pl. vi fig. 6.—S. bathypetalus, brachypetalus, dumblei, Roig 170 p. 175.—S. nummulitica, Shikama 182 pl. xxxii fig. 18.—S. iolti pp. 42, 94 pl. ii fig. 7, karreri pp. 41, 93 pl. ii fig. 5, sardiniensis pp. 42, 94 pl. ii fig. 6, Szörényi 214.

Schizaster (Ova) canaliferus, Tortonese 220 p. 223.

†Schizechinus duciei pp. 12, 58 pl. viii figs.' 1-2, hungaricus pp. 12, 59 pl. ii fig. 1, Szörényi 214.

†Scutella almerai parva subsp. n. pp. 19, 67 pl. i fig. 6, S. eichwoldi sp. n. pp. 19, 67 pl. i fig. 5, Miocene Ukraine, Szörényi Geol. hung. Ser. Palaeont. 23; S. cf. leognanensis pp. 16, 64 pl. i figs. 7, S. paulensis pp. 17, 65 pl. i figs. 1-3, pl. ii fig 2, vindobonensis pp. 16, 62 pl. i figs. 4, 8, P. sp. pp. 17, 65 pl. iii fig. 1, ID.—S., discussion of genus with type species S. subrotunda (neotype selected), Durham 50 p. 347 figs. 18, 1D, pl. 47 figs. 1-3.—Scutella patagonensis, Köster 93 p. 288 pl. i figs. 1-8.

†Scutellaster with cretaceous fig. 1B, interlineata fig. 1C, DURHAM 49.

†Sinaechinus kawaguchii a synonym of Faorina chinensis, HAYASAKA 71 p. 218.

†Sismondia naganoensis sp. n. Miocene Japan, Morishita Mem. Coll. Sci. Kyoto B 20 pp. 218–219 fig. 2, pl. i figs. 1-2.—S. convexa, Shikama 182 pl. xxxii fig. 19.

†Spatangomorpha see Eupatagus.

Spatangus californicus, H. L. CLARK 30 p. 345 fig. 3, pl. lxix figs. 78-79.—S. purpureus, larval stages, Rees 165 figs.—S. inermis p. 221, purpureus p. 220, TORTONESE 220.—S. purpureus, Turbellaria parasitic in, Westelad 238.

Sphaerechinus granularis, Palombi & Santarelli 155 p. 214 figs.—S. granularis, Tortonese 220 p. 215.—S. granularis, Ubisch 227 p. 294 fig. 3 [Pluteus].—S. granularis, hybrids with Paracentrotus lividus figs. 2, 4, and Psammechinus microtuberculatus, Ubisch 227 p. 294 [Plutei].

Sterechinus agassizii p. 18 pl. ix figs. 1-3, pl. x figs. 1-6, antarcticus p. 22, neumayeri p. 21 pl. xx fig. 4, Bebnasoni 8.

†Stigmatopygus malheiroi, DARTE-VELLE 41 p. 52.

Stolonoclypus see Clypeaster.

Strongylocentrotus franciscanus p. 278 pl. xlv fig. 23, purpuratus p. 279 pl. xliv fig. 22, H. L. Clark 30.—S. purpuratus, IRWIN 30 p. 307.—S. droebachiensis, echinoides, franciscanus, purpuratus, SWAN 210.

†Studeria rositae sp. n. Oligocene Cuba, Roig An. Acad. Habana 91 p. 147 pl. iv figs. 6-7.

Stylocidaris affinis, Tortonese 220 p. 211.

†Tetragramma bösei p. 30, malbosii p. 31, streeruwitzi p. 31 pl. ii fig. 21, T. sp. p. 32 pl. ii figs. 32–34, Maldo-NADO-KOERDELL 115.

Tetrapygus niger, H. L. Clark 30 p. 247 pl. xl fig. 13.

Tholeopelta see Clypeaster.

†Thylechinus sp. ?, DARTEVELLE 41 p. 23 pl. iii figs. 1-2, 16.

†Toxaster aff. granosus var. holasteroides, T. subcavatus, Gigout 63 p. 351.

Toxopneustes roseus, H. L. CLARK 30 p. 262 pl. xlii fig. 17.

Tripneustes depressus H. L. Clark 30 p. 263 pl. xlii fig. 18.—T. gratilla, Dakin, Bennett & Pope 40 p. 331 pl. xcii fig. 3.—T. esculentus, Fontaine 60 p. 5 figs.

†Tripylaster philippii, Bernascont 8 p. 53 pl. xxvii figs. 3-4, 6, pl. xxvii fig. 5, pl. xxix figs. 4-6, pl. xxx, pl. xxxi figs. 1-9.

Tripylus excavatus, Bernasconi 8 p. 51 pl. xxii figs. 1-6.

†Tristomanthus podjarkóvi pp. 25, 74 pl. iv fig. 4, podolicus pp. 26, 75 pl. vi fig. 6, spp. n. Miocene Ukraine, Szörényi Geol. hung. Ser. Palseont. 23; angulosus pp. 22, 71 pl. iii fig. 6, aremoricus pp. 23, 72 pl. iii fig. 4, fischeuri pp. 24, 74 pl. vi fig. 5, meslei pp. 24, 73 pl. iv fig. 3, subcylindricus pp. 23, 71 pl. v fig. 4, ID.

†SOMASTEROIDEA

[Vacant.]

ASTEROIDEA

Allostichaster polyplax, Dakin, Bennett & Pope 40 p. 319 pl. lxxix fig. 3.— A. insignis, Fell 55 p. 94.

Anseropoda placenta, Tortonese 220 p. 194.

†Asteracites quinquefolius, Sehlacher 180 p. 102 fig. la, pl. x.

†Asterias spp., MULLER 148 p. 23 figs. 7, X-Z, AA-BB.

Asterina aucklandensis, FELL 55 p. 88.

—A. folium p. 183 fig., hartmeyeri p. 183, FONTAINE 58.—A. gibbosa p. 192, pancerii p. 194 pl. i figs. 5–6, TORTONESE 220.

Asterodon robustus sp. n. Subantaretie, Fell Dom. Mus. Rec. Zool., Wellington 2 pp. 80–87 fig. l, pl. i figs. A–C, pl. ii figs. C–D; dilatatus p. 77 pl. ii figs. A–B, miliaris p. 77 pl. ii figs. E–F, ID.

Astropecten pectinatus p. 321 pl. lxxxi fig. 3, polyacanthus p. 321 pl. lxxxi fig. 4, Dakin, Bennett & Pope 40.—
A. articulatus p. 180, duplicatus p. 181 fig., Fontaine 58.—A. aranciacus p. 182, bispinosus p. 182, irregularis pentacanthus p. 172 pl. i figs. 1—4, jonstoni p. 181, platyacanthus p. 183, spinulosus p. 184, Tortonese 220.

†Astropecten? pygmaeus, MÜLLER 148 p. 63 figs. AB, BA.

†Calliderma atagensis, BATALLER 6 p. 69 figs.

Calvasterias laevigata p. 96, suteri p. 98, Fell **55**.

Ceramaster placenta, TORTONESE 220 p. 187 fig. 3.

Chaetaster longipes, TORTONESE 220 p. 187.

†Chomataster acules p. 59 fig. QQ, spenceri p. 60 fig. CA, MÜLLER 148.

Chunaster scapanephorus, MADSEN 111 p. 80.

Coscinasterias calamaria, Dakin, Bennett & Pope 40 p. 318 pl. lxxix figs. 1-2, pl. lxxx fig. 2.—C. tenuispina, Tartarin 215 p. 1 figs. 8-16, 19-57.—C. tenuispina, Tortonese 220 p. 195.

Echinaster echinophorus, FONTAINE 58 p. 183 fig.—E. sepositus, TORTONESE 220 p. 191.

Eremicaster pacificus, MADSEN 111 p. 76 fig. 1.

Freyella benthophila, sexradiata, tuberculata, MADSEN 111 p. 84.

Hacelia attenuata, TORTONESE 220 p. 190 fig. 4.

Henricia compacta p. 91, lukinsii p. 90, H. sp. p. 93, Fell 55.

Hippasteria colossa p. 411, derjugini p. 412, leiopelta with subsp. aculeata and longimana p. 413, H. mammifera p. 409, pedicellaris p. 410, DJAKONOV 52.

Hyphalaster parfaiti, Madsen 111 p. 82.

Linckia guildingii, [Anon.] 1 p. 188 figs. 1-5.—L. guildingii, Fontaine 58 p. 182 fig.

Luidia maculata forma herdmani f.n. Indo-Pacific, Clark Bull. Brit. Mus. (nat. Hist.) Zool. 1 pp. 389-390 fig. 8, pl. xli figs. 2-3; L. africana p. 393 figs. 10-l1, alternata numidica p. 388 pl. xli fig. 1, L. aspera p. 386 figs. 4, 6, columbia p. 381 figs. 1-2, pl. xxxix fig. 1, forficifer p. 391, hardwicki p. 391 pl. xxxix figs. 2-3, magnifica p. 386 figs. 5, 7, mauritiensis p. 386, quinaria p. 392.

fig. 9, sarsi p. 393 fig. 12, savignyi p. 385 pl. xl fig. 2, scotti p. 383 fig. 3, pl. xl fig. 1, ID.—L. australiae, Dakin, Bennett & Pope 40 p. 321 pl. lxxxii.—L. alternata p. 181, clathrata p. 181 fig., senegalensis p. 182 fig., Fontaine 58.—L. sarsi, Tortonese 219 p. 269.—L. ciliaris p. 185, sarsi p. 186, Tortonese 220.

Luidiaster dawsoni ochotensis, DJA-KONOV 52 p. 409.

Marthasterias glacialis, TORTONESE 220 p. 196 fig. 5.

†Metopaster, Bataller 6 p. 68 figs.— M. poulsenii p. 53 figs. 5, 8, C, N, MM, XA-YA, DA, aff. poulsenii p. 54 figs. 2, B, J, M, V, JJ-LL, JA-KA, tumidus p. 52 figs. D, K, R-S, NN-PP, HA, LA, TA, UA, undulatus p. 55 figs. TT-WW, pl. xi fig. 3, Müller 148.

Ophidiaster guildingii, Fontaine 58 p. 182.—O. ophidianus, Tortonese 220 p. 189.

†Ophryaster magnus, Müller 148 p. 60 figs. F, ? H, ? O.

Oreaster reticulatus, Fontaine 58 p. 182 fig.—O. nodosus, Ohshima 153 p. 158 fig. 1, pl. i figs. 1—3.

Patiriella calcar p. 317 pl. lxxx fig. 1, pl. lxxxiii, exigua p. 317 pl. lxxx fig. 3, gunnii p. 318 pl. lxxx fig. 4, Dakin, Bennett & Pope 40.

Pentagonaster, diagnosis p. 398, abnormalis p. 398, dubeni p. 400 fig. 13b, pls. xliii-xliv, gunni p. 400, pulchellus p. 398 fig. 13a, pl. xlii, Clark 28.

Petricia vernicina, Dakin, Bennett & Pope 40 p. 312 pl. lxxxi fig. 5.

Plectaster decanus, Dakin, Bennett & Pope 40 p. 320 pl. lxxxi fig. 1.

Porcellanaster caeruleus, Madsen 111 p. 74.

†Pycinaster rasmusseni sp. n. Cretaceous Rügen, MÜLLER Geologia Beihft 8 pp. 49-51 figs. A, G, L, U, DD-HH, EA-GA, ZA-BC.

†Recurvaster radiatus, MÜLLER 148 p. 56 figs. 6, P, Q, T, XX-ZZ, VA-WA.

Sclerasterias richardi, Tortonese 219 p. 269.—S. richardi, Tortonese 220 p. 201 fig. 6, pl. i fig. 7.

Stegnaster wesseli, Fontaine 58 p. 183.

Stichaster australis, Fell 55 p. 93.

Styracaster horridus, Madsen 111 p. 78 fig. 2.

†Teichaster favosus, Müller 148 p. 58 figs. RR-SS.

Tethyaster subinermis, Tortonese 220 p. 185.

Thoracaster cylindratus, Madsen 111 p. 83.

Tosia, diagnosis p. 403, auranta p. 408, australis with forma astrologorum p. 404 fig. 14, pl. xlv figs. 1-2, pl. xlv fig. 3, T. grandis p. 408, magnifica p. 408 pl. xlv figs. 5, pl. xlvi figs. 4-5, nobilis p. 406 pl. xlv figs. 3-4, 6-7, pl. xlvi figs. 1-2, queenslandensis p. 411, rubra p. 406, tubercularis p. 406, Clark 28.

Uniophora granifera, Dakin, Bennett & Pope 40 p. 320 pl. lxxxi fig. 2.

†Valettaster ocellatus, MÜLLER 148 p. 61 figs. E, MA.

OPHIUROIDEA

Amphiodia repens, Fontaine 59 p. 201.

Amphiophiura bullata, MADSEN 112 p. 107 fig. 1.

Amphioplus basilicus, Fell 55 p. 102.

Amphipholis squamata, Fell 55 p. 102.—A. squamata, Fontaine 59 p. 201.—A. squamata, Tortonese 220 p. 208.

Amphiura amokurae, magellanica, Fell 55 p. 101.—A. stimpsonii, Fon-Taine 59 p. 200.—A. chiajei, filiformis, Tortonese 220 p. 207.

†Asteracites lumbricalis, SEILACHER 180 p. 94 figs. 2-3, pls. vii-viii.

Astrophyton muricatum, Fontaine 59 p. 200 fig.

Astrospartus mediterraneus, Torto-NESE 220 p. 203.

†Geocoma magna sp. n. Jurassic Spain, BATALLER An. Asoc. esp. Prog. Cienc. 6 1941 pp. 65-68 figs. 1-5. Ophiacantha setosa, Tortonese 219 p. 27; 220 p. 204.

Ophiactis savignyi, Dakin, Bennett & Pope 40 pl. lxxxvi upper fig.—O. savignyi, Fontaine 59 p. 201 fig.

Ophiarachnella ramsayi, Dakin, Bennett & Pope 40 p. 325 pl. lxxxvi lower fig., pl. lxxxvii.

Ophioceres marginata sp. n. Subantarctic, Fell Dom. Mus. Rec. Zool., Wellington 2 pp. 103-105 fig. 5, pl. i figs. D-E.

Ophiocnida scabriuscula, Fontaine 59 p. 201.

Ophiocoma pulchra, Dakin, Bennett & Pope 40 p. 324 pl. lxxxv lower fig.— O. echinata p. 203, pumila p. 203, riisei p. 203 fig., Fontaine 59.

Ophioderma appressum, brevicaudum, brevispinum, cinereum p. 204 figs., guttatum, rubicundum p. 205, Fontaine 59.—O. longicaudum, Tortonese 220 p. 208.

Ophiolepis elegans, paucispina, Fon-TAINE 59 p. 205.

Ophiomyxa brevirima, Fell 55 p. 100.—O. flaccida, Fontaine 59 p. 200 fla.—O. pentagona, Tortonese 220 p. 203.

Ophionereis, discussion and key to spp. with O. amoyensis sp. n. S. China pp. 88-90 fig. 11, intermedia sp. n. N. Japan pp. 90-92 fig. 12, saeakii sp. n. Japan pp. 75-77 fig. 4, pl. iii figs. 1-2, Clark Proc. 2001. Soc. London 123; albomaculata p. 74, annulata p. 74 fig. 3b, pl. i figs. 3-4, dolabriformis p. 70, dubia p. 83 fig. 9, pl. ii figs. 1-2 with subsp. sinensis p. 83 fig. 10, fusca p. 78 figs. 5-6, porrecta p. 80, reticulata p. 73 fig. 3a, pl. i figs. 1-2, semoni p. 82 fig. 8, pl. ii figs. 3-4, serradia p. 71, squamulosa p. 71 fig. 2, pl. iii figs. 3-4, variegata p. 81 fig. 7, Id. -O. schayeri, Dakin, Bennett & Popfe 40 p. 325 pl. lixxiv. O. reticulata, MILLOTT 124 p. 96 figs. 1-2.

Ophiosphalma armigerum p. 109, planum p. 108 fig. 2, Madsen 112.

Ophiostigma isacanthum, Fontaine 59 p. 201.

Ophiothrix angulata p. 202 fig., orstedii p. 202, suensonii p. 202, Fon-TAINE 59.—O. fragilis p. 204, quinquemaculata p. 206, TORTONESE 220.

Ophiozona impressa, Fontaine 59 p. 205.

Ophiura albida p. 210, texturata p. 209, Tortonese 220.

Pectinura gracilis, FELL 55 p. 102.

Placophiothrix spongicola, Dakin, Bennett & Pope 40 p. 325 pl. lxxxv upper fig.

CRINOIDEA

†Indet., Chronic 27 p. 165 pl. xxxv figs. 7–12.—Indet., Valensi 229 p. 69 pl. xii fig. 7.—Plate undet., Walter 235 pl. lxxiii fig. 8.

†Columnals, classification of, VYALOV 234.

†Abactinocrinus gen. n. for type A. rossei sp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27 pp. 528-530 pl. liii figs. 8-10, pl. lv figs. 22-23.

†Actinocrinites sp., Laudon & Severson 101 p. 528 pl. liii fig. 7.

†Actinocrinus higuchisawensis pp. 358–359 pl. i fig. 16, ohmoriensis pp. 359–360 pl. iv fig. 5, spp. n. Carboniferous Japan, Minato J. Fac. Sci. Hokkaido Univ. Geol. Min. 7 1951.—A. higuchisawaensis, Minato 128 p. 283 pl. i fig. 8.

†Amphelecrinus madisonensis pp. 521-522 pl. li figs. 20-21, pl. lv figs. 10-11, sacculatus p. 522 pl. li fig. 1, pl. lv fig. 7, spp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27.

†Amphoracrinus sp., Minato 130 p. 360.

Antedon mediterranea, TORTONESE 220 p. 168 fig. 2.

†Aorocrinus douglassi, Laudon & Severson 101 p. 527 pl. liii figs. 11-12.

†Archaeocrinus subovalis sp. n. Ordovician U.S.A., STRIMPLE J. Paleont. 27 p. 606 figs. 1-7.

†Austinocrinus with Austinocrinus subgen. n. for type Austinocrinus komarolfi de Loriol 1889 Cretaceous Turkestan, Sieverns-Doreck Mitt. geol. (St.) Inst. Hamb. 22 p. 113; rothpletzi p. 103 figs. 1-6, pl. xviii figs. 1-13, pl. xviii figs. 1-7, 12, rothpletzi var. p. 112 pl. xviii fig. 14, pl. xviii figs. 8-11, Id.

†Austinocrinus with Penroseocrinus subgen, n. for type Balanocrinus mexicanus Springer 1922 Cretaceous Mexico, Sievers-Doreck Mitt. geol. (St.) Inst. Hamb. 22 p. 114 pl. xviii fig. 13.

†Aviadocrinus with A. sampelayoi, Sieverts-Doreck 186 p. 85 pl. iv fig. 11.

†Bactrocrinus, Sieverts-Doreck 186 p. 79.

†Batocrinus gallatinensis sp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27 pp. 526–527 pl. lii figs. 23–27, pl. lv figs. 19–21.

†Bridgerocrinus gen. n. for type B. fairgensis sp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27 pp. 517-519 pl. li figs. 6-8, pl. lv fig. 5.

†Cactocrinus magnidactylus sp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27 pp. 531–532 pl. liii fig. 5; arnoldi p. 530 pl. liii figs. 1–3, nodobrachiatus p. 530 pl. liii figs. 6, ornatissimus p. 530 pl. liii fig. 4, In.

†Calceocrinid, Sieverts-Doreck 186 p. 78 fig. 8.

†Carinocrinus eventus sp. n. Carboniferous U.S.A., STRIMPLE J. Wash. Acad. Sci. 43 pp. 201–203 figs. 1–2.

Cenolia trichoptera, Dakin, Bennett & Pope 40 p. 327 pl. lxxxviii.

†Cradeocrinus dendratus sp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27 p. 514 pl. li figs. 9–12, pl. lv figs. 3–4.

†Cribanocrinus watersianus, Laudon & Severson 101 p. 526 pl. lii figs. 20–22.

†Crotalocrinus? sp., HOLTEDAHL 75 p. 203 fig. 78, 18 [stem].

†Culmicrinus jeffersonensis sp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27 p. 515 pl. li figs. 29-30, pl. lv fig. 1.

†Cupressocrinus with C. abbreviatus figs. 6-7, pl. iv fig. 10, crassus figs. 1, 3-4, pl. iv figs. 8-9, C.? gibber fig. 5, C. gracilis pl. iv fig. 7, C. sp. fig. 2, SIEVERTS-DORECK 186 p. 82.

†Cyathocrinites sp., Laudon & Severson 101 p. 513 pl. li figs. 13-14.

†Dichocrinus quadriceptatus sp. n. Carboniferous U.S.A., Laudon & Severson J. Paleont. 27 pp. 534–535 pl. liv figs. 11–14, pl. lv fig. 26; bozemanensis p. 533 pl. liv figs. 15–16, campto p. 534 pl. liv fig. 17, douglassi p. 535 pl. liv fig. 10, Id.

†Dinotocrinus logani sp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27 p. 516 pl. li figs. 23-27, pl. lv figs. 12-14.

†Encrinus liliiformis, KUHN-SCHNY-DER 96 p. 210 fig. 2b.

†Eutaxocrinus montanensis, LAUDON & SEVERSON 101 p. 523 pl. liv figs. 18-20.

†Goniocrinus sculptilis, Laudon & Severson 101 p. 514 pl. li fig. 22.

†Holcocrinus longicirrifer, LAUDON & SEVERSON 101 p. 522 pl. li figs. 31-35, pl. lv fig. 2.

†Hypselocrinus maccabei, Laudon & Severson 101 p. 519 pl. li figs. 2-5.

†Isocrinus, discussion of genus p. 122, nehalemensis sp. n. pp. 135-135 figs. 35, 36 B-C, 37-39 A-C, pl. xix figs. 1, pl. xx figs. 1-2, pl. xxi fig. 2, pl. xxii fig. 1, pl. xxiii figs. 4, 7, 11, oregonensis sp. n. pp. 124-135 figs. 29-35, 36A, 36 D-E, 39D, pls. xiv-xxiii, pl. xix figs. 2-5, pl. xx fig. 3, pl. xxii fig. 3, pl. xxiii figs. 1-3, 5-6, 8-10, pl. xxiv, Oligocene? U.S.A., Moore & Vokes Prof. Pap. U.S. Geol. Surv. 233E.

†Lecanocrinus with subgen. Miracrinus subgen. n. for type M. perdewi sp. n. Devonian U.S.A., Bowsher Smithson. misc. Coll. 121 9 pp. 3-7 figs. 1 a-e, pl. i figs. 1-11; L. (Lecanocrinus) macropetalus, Io. p. 2 figs. 1 f-g.

Leptometra phalangium, Tortonese 220 p. 171.

†Linocrinus wilsallensis sp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27 p. 517 pl. li fig. 28, pl. lv fig. 15.

†Miracrinus subgen. n. see Lecanocrinus.

†Myelodactylus canaliculatus p. 76 pl. iv figs. 1-3, M. sp. p. 78 pl. iv figs. 4-6, Sieverts-Doreck 186.

†Parulocrinus? sp., Chronic 27 p. 164 pl. xxxv figs. 5-6.

†Pelecocrinus sp., Laudon & Severson 101 p. 515 pl. li fig. 15, pl. lv fig. 6.

Penroseocrinus subgen. n., see Austinocrinus.

†Pentacrinus sp., MÜLLER 147 pl. i fig. 2.

†Pentagonocyclopa cf. dispar, Walter 235 p. 701 pl. lxiii figs. 15-17.

†Periechocrinites sp., Laudon & Severson 101 p. 527 pl. lii figs. 1-2.

†Platycrinites basicraniatus p. 533 pl. liv fig. 1, pl. lv fig. 24, canadensis pp. 532-533 pl. liv fig. 8, pl. lv fig. 25, spp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27; bozemanensis p. 532 pl. liv figs. 2-7, incomptus p. 532 pl. liv figs. 9, ID.

†Platycrinus asiatica sp. n. Carboniferous Japan, Minato J. Fac. Sci. Hokkaido Univ. Geol. Min. 7 1951 pp. 357–358 pl. i fig. 8, pl. iv figs. 2a-b.—P. asiatica, Minato 128 p. 287 pl. i fig. 1.

†Pseudosaccocoma japonica, Shikama 182 pl. xix fig. 8.

†Ramulocrinus livingstonensis sp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27 pp. 519-520 pl. li figs. 16-19, pl. lv figs. 8-9.

†Rhodocrinites macrotumidus pp. 524-525 pl. lii figs. 9-12, pl. lv figs. 17-18, priminodosus pp. 525-526 pl. lii figs. 6-8, pl. lv fig. 16, spp. n. Carboniferous U.S.A., LAUDON & SEVERSON J. Paleont. 27; cirrusi p. 524 pl. lii figs. 3-5, douglassi p. 524 pl. lii figs. 13-19, ID.

†Rhopalocrinus with R. gracilis, Sieverts-Doreck 186 p. 85 pl. iv fig. 12.

†Solanocrinus astrei pp. 61-63 figs. 1-3, estruchi pp. 63-65 figs. 4-12, spp. n. Cretaceous Spain, Bataller An. asoc. esp. Prog. Cienc. 6 1941.

Tropiometra carinata, Fontaine 61 p. 33.

†MACHAERIDIA

Lepidocoleus sularpensis sp. n. Ordovician Sweden, Lindström Geol. Fören. Stockh. Förh. 75 p. 143 pl. i fig. 7.

†EOCRINOIDEA

Gogia prolifica, HARKER & HUTCHIN-SON 67 p. 285 pl. xl figs. 1-5.

Lichenoides priscus, UBAGHS 223 p. 3 figs. 1-11.

†PARACRINOIDEA

[Vacant.]

†CYSTIDEA

Camarocrinus asiaticus, Sun & Szetu 209 p. 243 pl. i.

Echinosphaerites aurantium, Holte-DAHL 75 p. 165 fig. 74, 4-5.

†CARPOIDEA

Decacystis hispanicus, Melendez 121 p. 509 fig. 8.

Myeinocystites gen. n. p. 105 for type M. natus sp. n. Ordovician U.S.A., STRIMPLE J. Wash. Acad. Sci. 43 pp. 105-106 figs. 1-2.

Trochocystites bohemicus, MELENDEZ 121 p. 504 figs. 3-6, 8, pl. i figs. 1-6.

†EDRIOASTEROIDEA

Agelacrinites jasperensis sp. n. Carboniferous Canada, HARKER J. Paleont.

27 pp. 288–289 fig. 1.—A. taunicus sp. n. Devonian Germany, RÖSLER Notizblihess. Landessamt. Bodenforsch. 31 pp. 70–72 pl. iii figs. 1–3; ? rhenanus, In. p. 72.

†BLASTOIDEA

Gen. et sp. indet., Minato 128 p. 287 pl. i fig. 5.—Gen. et sp. indet., Minato 130 p. 361 pl. i figs. 5, 9, 12, pl. iv fig. 6. Codaster acutus, trilobatus, Joysey 86 p. 208 fig. 1.

Cryptoschisma schultzi, Bergou-NIOUX 7 p. 636 figs. 9-12.

INCERTAE SEDIS

Echinoderm fragments, VALENSI 229 p. 69 figs. 4 d-e.